

Turkish Journal of Chemistry

Turkish Journal

of

Chemistry

Terpenoids and Steroids from the Roots of *Salvia blepharochlaena*

Ufuk KOLAK¹, Gülaçtı TOPÇU¹, Seher BİRTEKSÖZ²,
Gülten ÖTÜK², Ayhan ULUBELEN¹

¹ İstanbul University, Pharmacy Faculty, Department of General Chemistry
34116, İstanbul-TURKEY

e-mail: ufukkolak@yahoo.com

²İstanbul University, Pharmacy Faculty, Department of Pharmaceutical Microbiology
34116, İstanbul-TURKEY

 [Keywords](#)
 [Authors](#)



chem@tubitak.gov.tr

[Scientific Journals Home](#)
[Page](#)

Abstract: From the roots of *Salvia blepharochlaena* Hedge and Hub. Mor. 4 triterpenoids, 4 steroids, 6 diterpenoids, and an aromatic ester were isolated. The structures of these compounds were established by spectroscopic methods. Formosanolide was isolated for the first time from the genus *Salvia*.

Key Words: *Salvia blepharochlaena* Hedge and Hub. Mor., Lamiaceae, triterpenoids, steroids, diterpenoids, aromatic ester

Turk. J. Chem., **29**, (2005), 177-186.

Full text: [pdf](#)

Other articles published in the same issue: [Turk. J. Chem., vol.29, iss.2.](#)