Turkish Journal of Chemistry

Turkish Journal

of

Chemistry

Chemical Constituents of Campanula lactiflora

Nurettin YAYLI, Nuri YILDIRIM, Asu USTA, Serpil ÖZKURT, Vildan AKGÜN Karadeniz Technical University, Faculty of Science, Department of Chemistry 61080 Trabzon-TURKEY e-mail: yayli@ktu.edu.tr

Keywords Authors <u>Abstract:</u> A new flavonoid glycoside, 4'-O-(p-hydroxybenzoyl)-isorhamnetin-3,7- di-O-β-D-glucopyranoside (1), and 4 known compounds, sitosterol β-D-glucoside (2), bis(2-ethylhexyl) adipate (3), p-hydroxybenzoic acid (4) and ethyl docosanoate (5), were isolated for the first time from Campanula lactiflora and their structures were deduced by high field 1D and 2D NMR and (+) FAB and EI mass spectra.



<u>Key Words:</u> Campanulaceae, Campanula lactiflora, flavonoid glycoside, steroid, aromatic, ester

chem@tubitak.gov.tr

Turk. J. Chem., 27, (2003), 749-756.

Full text: pdf

Scientific Journals Home Page Other articles published in the same issue: Turk. J. Chem., vol. 27, iss. 6.