



# Turkish Journal of Chemistry

Turkish Journal

of

Chemistry

 [Keywords](#)  
 [Authors](#)



[chem@tubitak.gov.tr](mailto:chem@tubitak.gov.tr)

[Scientific Journals Home Page](#)

## Secondary metabolites from *Nepeta heliotropifolia*

Zühal GÜVENALP<sup>1</sup>, Hilal ÖZBEK<sup>1</sup>, Ayşe KURUÜZÜM-UZ<sup>2</sup>,  
Cavit KAZAZ<sup>3</sup>, L. Ömür DEMİREZER<sup>2</sup>

<sup>1</sup>Atatürk University, Faculty of Pharmacy, Department of Pharmacognosy,  
TR-25240 Erzurum-TURKEY  
e-mail: zguvenalp@atauni.edu.tr

<sup>2</sup>Hacettepe University, Faculty of Pharmacy, Department of Pharmacognosy,  
TR-06100 Ankara-TURKEY

<sup>3</sup>Atatürk University, Faculty of Arts and Sciences, Department of Chemistry,  
TR-25240 Erzurum-TURKEY

**Abstract:** Two iridoid glycosides, ixoroside (1) and nepetanudoside B (2); 1 phenylpropanoid glycoside, coniferine (3); 2 flavone glycosides, apigenin 7-O-glucuronide (4) and apigenin 7-O-glucopyranoside (5); 2 triterpenes, oleanolic acid (6) and ursolic acid (7); and 1 sterol,  $\beta$ -sitosterol (8), were isolated from the aerial parts of *Nepeta heliotropifolia* Lam. (Lamiaceae). Their structures were identified by means of spectroscopic methods (1D- and 2D-NMR, UV, and EIMS).

**Key Words:** Lamiaceae, *Nepeta heliotropifolia*, iridoid glycosides, flavone glycosides, triterpenoids.

---

Turk. J. Chem., **33**, (2009), 667-675.

Full text: [pdf](#)

Other articles published in the same issue: [Turk. J. Chem.,vol.33,iss.5.](#)