Turkish Journal of Chemistry

Turkish Journal

Calculation of Multi-Center Integrals for the Region of Moderate and Weak Overlap

of

Chemistry

Abdulhalik KARABULUT, Gökhan BUDAK, Lütfü DEMİR
Atatürk University, Sciences and Letters Faculty,
Department of Physics, 25240 Erzurum-TURKEY
Nagif Sefteroğlu NABİYEV
Bakü State University, Faculty of Physics, Bakü-AZERBAJCAN
Yusuf ŞAHİN
Atatürk University, Sciences and Letters Faculty,

Atatürk University, Sciences and Letters Faculty, Department of Physics, 25240 Erzurum-TURKEY

Keywords Authors



chem@tubitak.gov.tr

Scientific Journals Home Page sets using the two-center expansion of nuclear attraction and electron repulsion potentials. The terms of the formula are the analytical functions of interatomic distances. Three kinds of integrals contained in these formulae can be evaluated analytically. In the presented method, unlike many other popular methods, the numerical integrations procedure is unnecessary since it does not contain the incomplete gamma function. A computer program was written and tested for various orbitals. The calculations show that the series converge very rapidly.

Abstract: In this study a general formula was established for multi-center integrals over Gaussian basis

Turk. J. Chem., 25, (2001), 405-418.

Full text: pdf

Other articles published in the same issue: Turk. J. Chem., vol. 25, iss. 4.