

# Turkish Journal of Chemistry

Turkish Journal

of

Chemistry

 [Keywords](#)  
 [Authors](#)



[chem@tubitak.gov.tr](mailto:chem@tubitak.gov.tr)

[Scientific Journals Home Page](#)

## Iridoid Glucosides from *Lamium garganicum* subsp. *laevigatum*

Tayfun ERSÖZ<sup>1</sup>, Duygu KAYA<sup>1</sup>, Funda Nuray YALÇIN<sup>1,2</sup>, Cavit KAZAZ<sup>3</sup>, Erhan PALASKA<sup>4</sup>, Charlotte H. GOTFREDSEN<sup>5</sup>, Søren R. JENSEN<sup>5</sup>, İhsan ÇALIŞ<sup>1</sup>

<sup>1</sup>Hacettepe University, Faculty of Pharmacy, Department of Pharmacognosy, TR-06100 Ankara, TURKEY

e-mail: tersoz@hacettepe.edu.tr

<sup>2</sup>Hacettepe University, Faculty of Pharmacy, Department of Pharmacy Management, TR-06100 Ankara-TURKEY

<sup>3</sup>Atatürk University, Faculty of Arts and Science, Department of Chemistry, TR-25240 Erzurum-TURKEY

<sup>4</sup>Hacettepe University, Faculty of Pharmacy, Department of Pharmaceutical Chemistry, TR-06100 Ankara-TURKEY

<sup>5</sup>The Technical University of Denmark, Department of Chemistry Building 201, DK-2800 Lyngby-DENMARK

**Abstract:** Phytochemical investigations on the above-ground parts of *Lamium garganicum* subsp. *laevigatum* resulted in the isolation of 7 iridoid glucosides: shanzhiside methyl ester (1), barlerin (8-O-acetylshanzhiside methyl ester) (2), 6-O-syringyl-8-O-acetylshanzhiside methyl ester (3), 6 $\beta$ -hydroxyipolamiide (4), lamalbide (5), dehydropenstemoside (6), and sesamoside (7). The structure of the iridoids was elucidated by spectroscopic (UV, IR, 1D- and 2D-NMR, and ESI-MS) evidence.

**Key Words:** *Lamium garganicum* L. subsp. *laevigatum*, Lamiaceae, iridoid glucosides

---

Turk. J. Chem., **31**, (2007), 155-162.

Full text: [pdf](#)

Other articles published in the same issue: [Turk. J. Chem., vol.31, iss.2.](#)