Astrophysics > Cosmology and Extragalactic Astrophysics

Luke-warm dark matter: Bosecondensation of ultra-light particles

Andrew P. Lundgren, Mihai Bondarescu, Ruxandra Bondarescu, Jayashree Balakrishna

(Submitted on 31 Dec 2009)

We discuss the thermal evolution and Bose condensation of ultra-light scalar particles with Compton wavelength of galactic scales. We find an upper bound of 1.5 K for the dark matter temperature from the WMAP constraints on the amount of hot dark matter for a \$\Lambda\$CDM model. Agglomerations of these particles can form stable halos and naturally prohibit small scale structure, which may be favored by observations of dark matter distributions near the centers of galaxies. We present numerical as well as approximate analytical solutions of the Friedmann-Klein-Gordon equations and study the cosmological evolution of this scalar field dark matter from the early universe to the era of matter domination. Today, the particles in the ground state mimic presureless matter, while the excited state particles are radiation like.

Comments: 5 pages, 2 figures, Submitted to ApJ Letters

Subjects: Cosmology and Extragalactic Astrophysics (astro-ph.CO); General Relativity and Quantum Cosmology (gr-qc) Cite as: arXiv:1001.0051v1 [astro-ph.CO]

Submission history

From: Ruxandra Bondarescu [view email] [v1] Thu, 31 Dec 2009 00:08:01 GMT (39kb)

Which authors of this paper are endorsers?

Link back to: arXiv, form interface, contact.

(Help | Advanced search)

All papers 🚽 Go!

Download:

- PostScript
- PDF
- Other formats

Current browse context: astro-ph.CO < prev | next > new | recent | 1001

Change to browse by:

astro-ph gr-qc

References & Citations

- SLAC-SPIRES HEP (refers to | cited by)
- NASA ADS
- CiteBase

Bookmark(what is this?)