



High Energy Physics - Theory

# Safety of Minkowski Vacuum

Gia Dvali

(Submitted on 5 Jul 2011)

We give a simple argument suggesting that in a consistent quantum field theory tunneling from Minkowski to a lower energy vacuum must be impossible. Theories that allow for such a tunneling also allow for localized states of negative mass, and therefore, should be inconsistent.

Comments: 3 pages, Latex

Subjects: **High Energy Physics - Theory (hep-th)**; Cosmology and Extragalactic Astrophysics (astro-ph.CO); General Relativity and Quantum Cosmology (gr-qc); High Energy Physics - Phenomenology (hep-ph)

Cite as: **arXiv:1107.0956 [hep-th]**  
(or **arXiv:1107.0956v1 [hep-th]** for this version)

## Submission history

From: Georgi Dvali [[view email](#)]

[v1] Tue, 5 Jul 2011 19:32:30 GMT (5kb)

*Which authors of this paper are endorsers?*

Link back to: [arXiv](#), [form interface](#), [contact](#).

## Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

## Current browse context:

hep-th

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1107](#)

## Change to browse by:

[astro-ph](#)  
[astro-ph.CO](#)

[gr-qc](#)  
[hep-ph](#)

## References & Citations

- [INSPIRE HEP](#)  
([refers to](#) | [cited by](#))
- [NASA ADS](#)

## Bookmark (what is this?)

