



Diffuse Neutrino Flux

[Jürgen Brunner](#)

(Submitted on 8 Jul 2011)

The search for a diffuse neutrino flux component from astrophysical sources complements searches for point sources. In 2010 and 2011 many new results have been published on this subject. Realistic models can now be tested with these new measurements from the leading neutrino telescope experiments. An overview over these recent results is given.

Comments: Contribution to Conference Proceedings of XIV International Workshop on "Neutrino Telescopes", Venice 2011

Subjects: **High Energy Astrophysical Phenomena (astro-ph.HE)**; High Energy Physics - Experiment (hep-ex)

Cite as: [arXiv:1107.1593](#) [astro-ph.HE]
(or [arXiv:1107.1593v1](#) [astro-ph.HE] for this version)

Submission history

From: Jürgen Brunner [[view email](#)]

[v1] Fri, 8 Jul 2011 09:55:27 GMT (42kb)

[Which authors of this paper are endorsers?](#)

Link back to: [arXiv](#), [form interface](#), [contact](#).

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

astro-ph.HE

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1107](#)

Change to browse by:

[astro-ph](#)

[hep-ex](#)

References & Citations

- [INSPIRE HEP](#)
([refers to](#) | [cited by](#))
- [NASA ADS](#)

Bookmark([what is this?](#))

