

# Large Extra Dimensions and Small Black Holes at the LHC

Marcus Bleicher, Piero Nicolini

(Submitted on 13 Jan 2010 (v1), last revised 24 Jan 2010 (this version, v3))

We present an overview on the conjectured production of microscopic black holes as a consequence of high energy hadronic collisions at the LHC, CERN (Geneva, Switzerland) from this year on. Provided the presence of large extradimensions, we analyze some possible scenarios of what could turn to be an outstanding experimental discovery. We also discuss some new models which have been recently proposed on the ground of quantum gravity arguments. The final comments are devoted to supposed potential risks connected to the formation of black holes in particle detectors.

Comments: 12 pages, 4 figures, 2 tables, Proceedings of 'Symmetries in Science', July 19-24, 2009 at Kloster Mehrerau, Bregenz, Austria - Updated reference list and modified conclusion

Subjects: **High Energy Physics - Phenomenology (hep-ph)**; General Relativity and Quantum Cosmology (gr-qc); High Energy Physics - Experiment (hep-ex); High Energy Physics - Theory (hep-th)

Cite as: **arXiv:1001.2211v3 [hep-ph]**

## Submission history

From: Piero Nicolini [[view email](#)]

[v1] Wed, 13 Jan 2010 15:48:10 GMT (49kb)

[v2] Fri, 22 Jan 2010 15:15:01 GMT (50kb)

[v3] Sun, 24 Jan 2010 21:51:52 GMT (50kb)

*[Which authors of this paper are endorsers?](#)*