



AMBER high-spectral resolution interferometry of Be stars: More than just stars and disks?

S. Stefl, T. Rivinius, D. Baade, A. Carciofi

(Submitted on 26 Jul 2011)

High spectral-resolution Br-gamma VLTI/AMBER observations of classical Be stars reveal complex visibility, phase, and closure phase profiles, which were not seen in previous lower-resolution data. They present new challenges for the modeling of Be stars and may require additional ingredients over and above the conventional central star + circumstellar disk paradigm.

Comments: Presented at the workshop "Resolving the Future of Astronomy with Long-Baseline Interferometry", Socorro, NM, USA, March 28 - 31, 2011

Subjects: **Solar and Stellar Astrophysics (astro-ph.SR)**

Cite as: [arXiv:1107.5297](#) [astro-ph.SR]

(or [arXiv:1107.5297v1](#) [astro-ph.SR] for this version)

Submission history

From: Stan Stefl [[view email](#)]

[v1] Tue, 26 Jul 2011 19:21:00 GMT (41kb)

[Which authors of this paper are endorsers?](#)

Link back to: [arXiv](#), [form interface](#), [contact](#).

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

astro-ph.SR

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1107](#)

Change to browse by:

[astro-ph](#)

References & Citations

- [INSPIRE HEP](#)
([refers to](#) | [cited by](#))
- [NASA ADS](#)

Bookmark([what is this?](#))

