

Home

Online Library

- Recent Papers
- Volumes
- Library Search
- Title and Author Search

RSS Feeds

General Information

Submission

Review

Production

Subscription

Journal Metrics

 not applicable

SCOPUS[®] SNIP 0.287

SCOPUS[®] SJR 0.054

Definitions

ARCHIVED IN


PORTICO

Volumes Contents of Volume 25

Adv. Geosci., 25, 29-36, 2010
www.adv-geosci.net/25/29/2010/
doi: 10.5194/adgeo-25-29-2010

© Author(s) 2010. This work is distributed
under the Creative Commons Attribution 3.0 License.

The forecaster's added value in QPF

M. Turco^{1,*} and M. Milelli¹

¹ARPA Piemonte (Regional Environmental Protection Agency), Torino, Italy
*now at: GAMA (Meteorological Hazards Analysis Team), Department of
Astronomy & Meteorology, Faculty of Physics, University of Barcelona, Barcelona,
Spain

Abstract. To the authors' knowledge there are relatively few studies that try to answer this question: "Are humans able to add value to computer-generated forecasts and warnings?". Moreover, the answers are not always positive. In particular some postprocessing method is competitive or superior to human forecast. Within the alert system of ARPA Piemonte it is possible to study in an objective manner if the human forecaster is able to add value with respect to computer-generated forecasts. Every day the meteorology group of the Centro Funzionale of Regione Piemonte produces the HQPF (Human Quantitative Precipitation Forecast) in terms of an areal average and maximum value for each of the 13 warning areas, which have been created according to meteo-hydrological criteria. This allows the decision makers to produce an evaluation of the expected effects by comparing these HQPFs with predefined rainfall thresholds. Another important ingredient in this study is the very dense non-GTS (Global Telecommunication System) network of rain gauges available that makes possible a high resolution verification. In this work we compare the performances of the latest three years of QPF derived from the meteorological models COSMO-17 (the Italian version of the COSMO Model, a mesoscale model developed in the framework of the COSMO Consortium) and IFS (the ECMWF global model) with the HQPF. In this analysis it is possible to introduce the hypothesis test developed by Hamill (1999), in which a confidence interval is calculated with the bootstrap method in order to establish the real difference between the skill scores of two competitive forecasts. It is important to underline that the conclusions refer to the analysis of the Piemonte operational alert system, so they cannot be directly taken as universally true. But we think that some of the main lessons that can be derived from this study could be useful for the meteorological community. In details, the main conclusions are the following:

- despite the overall improvement in global scale and the fact that the resolution of the limited area models has increased considerably over recent years, the QPF produced by the meteorological models involved in this study has not improved enough to allow its direct use: the subjective HQPF continues to offer the best performance for the period +24 h/+48 h (i.e. the warning period in the Piemonte system);

 Copernicus Publications
The Innovative Open Access Publisher

Search ADGEO

Full Text Search

Title Search

Author Search

News

- Please Note: Updated Reference Guidelines

Recent Papers

01 | ADGEO, 22 Nov 2010:
Tropopause and jetlet characteristics in relation to thunderstorm development over Cyprus

02 | ADGEO, 22 Nov 2010:
Probabilistic prediction of raw and BMA calibrated AEMET-SREPS: the 24 of January 2009 extreme wind event in Catalunya

03 | ADGEO, 15 Nov 2010:
Investigation of trends in synoptic patterns over Europe with artificial neural networks

– in the forecast process, the step where humans have the largest added value with respect to mathematical models, is the communication. In fact the human characterization and communication of the forecast uncertainty to end users cannot be replaced by any computer code;

– eventually, although there is no novelty in this study, we would like to show that the correct application of appropriated statistical techniques permits a better definition and quantification of the errors and, mostly important, allows a correct (unbiased) communication between forecasters and decision makers.

▣ [Full Article in PDF](#) (PDF, 633 KB)

Citation: Turco, M. and Milelli, M.: The forecaster's added value in QPF, Adv. Geosci., 25, 29-36, doi:10.5194/adgeo-25-29-2010, 2010. ▣ [Bibtex](#) ▣ [EndNote](#) ▣ [Reference Manager](#) ▣ [XML](#)