

[Home](#)[Online Library ACP](#)

- Recent Final Revised Papers
- [Volumes and Issues](#)
- Special Issues
- Library Search
- Title and Author Search

[Online Library ACPD](#)[Alerts & RSS Feeds](#)[General Information](#)[Submission](#)[Review](#)[Production](#)[Subscription](#)[Comment on a Paper](#)

Impact
Factor
4.865

ISI
indexed

[Volumes and Issues](#) [Contents of Issue 17](#)

Atmos. Chem. Phys., 8, 5279–5293, 2008

www.atmos-chem-phys.net/8/5279/2008/

© Author(s) 2008. This work is distributed under the Creative Commons Attribution 3.0 License.

Model simulations of stratospheric ozone loss caused by enhanced mesospheric NO_x during Arctic Winter 2003/2004

B. Vogel¹, P. Konopka¹, J.-U. Grooß¹, R. Müller¹, B. Funke², M. López-Puertas², T. Reddmann³, G. Stiller³, T. von Clarmann³, and M. Riese¹

¹ Forschungszentrum Jülich, ICG-1: Stratosphäre, Jülich, Germany

² Instituto de Astrofísica de Andalucía, CSIC, Granada, Spain

³ Forschungszentrum Karlsruhe, Institute for Meteorology and Climate Research, Karlsruhe, Germany

Abstract. Satellite observations show that the enormous solar proton events (SPEs) in October–November 2003 had significant effects on the composition of the stratosphere and mesosphere in the polar regions. After the October–November 2003 SPEs and in early 2004, significant enhancements of NO_x (=NO+NO₂) in the upper stratosphere and lower mesosphere in the Northern Hemisphere were observed by several satellite instruments. Here we present global full chemistry calculations performed with the CLaMS model to study the impact of mesospheric NO_x intrusions on Arctic polar ozone loss processes in the stratosphere. Several model simulations are performed with different upper boundary conditions for NO_x at 2000 K potential temperature (≈50 km altitude). In our study we focus on the impact of the non-local production of NO_x, which means the downward transport of enhanced NO_x from the mesosphere to the stratosphere. The local production of NO_x in the stratosphere is neglected. Our findings show that intrusions of mesospheric air into the stratosphere, transporting high burdens of NO_x, affect the composition of the Arctic polar region down to about 400 K (≈17–18 km). We compare our simulated NO_x and O₃ mixing ratios with satellite observations by ACE-FTS and MIPAS processed at IMK/IAA and derive an upper limit for the ozone loss caused by enhanced mesospheric NO_x. Our findings show that in the Arctic polar vortex (equivalent lat. >70° N) the accumulated column ozone loss between 350–2000 K potential temperature (≈14–50 km altitude) caused by the SPEs in October–November 2003 in the stratosphere is up to 3.3 DU with an upper limit of 5.5 DU until end of November. Further, we found that about 10 DU, but in any case lower than 18 DU, accumulated ozone loss additionally occurred until end of March 2004 caused by the transport of mesospheric NO_x-rich air in early 2004. The solar-proton-produced NO_x above 55 km due to the SPEs of October–November 2003 had a negligibly small impact on ozone loss processes through the end of November in the lower stratosphere (350–700 K ≈14–27 km). The mesospheric NO_x intrusions in early 2004 yielded a lower stratospheric ozone loss of about 3.5 DU, and clearly lower than 6.5 DU through the end of March. Overall, the non-local production of NO_x is an additional variability in the existing variations of the ozone loss observed in the Arctic.

[Search ACP](#)

Library Search

Author Search

[News](#)

- Sister Journals AMT & GMD
- Financial Support for Authors
- Journal Impact Factor
- Public Relations & Background Information

[Recent Papers](#)

01 | ACPD, 19 Nov 2008:
Stratospheric BrONO₂
observed by MIPAS

02 | ACPD, 19 Nov 2008:
Methyl chavicol:
characterization of its
biogenic emission rate,
abundance, and oxidation
products in the atmosphere

03 | ACP, 19 Nov 2008:
Technical Note: Quantitative
long-term measurements of
VOC concentrations by PTR-
MS – measurement,
calibration, and volume
mixing ratio calculation
methods

▣ [Final Revised Paper](#) (PDF, 5550 KB) ▣ [Supplement](#) (2231 KB) ▣ [Discussion Paper](#) (ACPD)

Citation: Vogel, B., Konopka, P., Grooß, J.-U., Müller, R., Funke, B., López-Puertas, M., Reddmann, T., Stiller, G., von Clarmann, T., and Riese, M.: Model simulations of stratospheric ozone loss caused by enhanced mesospheric NO_x during Arctic Winter 2003/2004, *Atmos. Chem. Phys.*, 8, 5279-5293, 2008. ▣ [Bibtex](#) ▣ [EndNote](#) ▣ [Reference Manager](#)