

[Home](#)[Online Library ACP](#)

- Recent Final Revised Papers
- [Volumes and Issues](#)
- Special Issues
- Library Search
- Title and Author Search

[Online Library ACPD](#)[Alerts & RSS Feeds](#)[General Information](#)[Submission](#)[Review](#)[Production](#)[Subscription](#)[Comment on a Paper](#)

Impact
Factor
4.865

ISI
indexed

[Volumes and Issues](#) [Contents of Issue 3](#)

Atmos. Chem. Phys., 5, 703-714, 2005
www.atmos-chem-phys.net/5/703/2005/
© Author(s) 2005. This work is licensed
under a Creative Commons License.

The role of organic aerosols in homogeneous ice formation

B. Kärcher¹ and T. Koop²

¹DLR Oberpfaffenhofen, Institute for Atmospheric Physics, Wessling, Germany

²University of Bielefeld, Faculty of Chemistry, Bielefeld, Germany

Abstract. Recent field observations suggest that the fraction of organic-containing aerosol particles in ice cloud particles is diminished when compared to the background aerosol prior to freezing. In this work, we use model calculations to investigate possible causes for the observed behavior. In particular, homogeneous freezing processes in cooling air parcels containing aqueous inorganic particles and organic particles are studied with a detailed microphysical model. A disparate water uptake and resulting size differences that occur between organic and inorganic particles prior to freezing are identified as the most likely reason for the poor partitioning of organic aerosols into the ice phase. The differences in water uptake can be caused by changes in the relationship between solute mass fraction and water activity of the supercooled liquid phase, by modifications of the accommodation coefficient for water molecules, or by a combination thereof. The behavior of peak ice saturation ratios and total ice crystal number concentrations is examined, and the dependence of the results on cooling rate is investigated. Finally, processes are discussed that could possibly modify the homogeneous freezing behavior of organic particles.

[Final Revised Paper](#) (PDF, 302 KB) [Discussion Paper](#) (ACPD)

Citation: Kärcher, B. and Koop, T.: The role of organic aerosols in homogeneous ice formation, Atmos. Chem. Phys., 5, 703-714, 2005. [Bibtex](#) [EndNote](#) [Reference Manager](#)

[Search ACP](#)

Library Search

Author Search

[News](#)

- Sister Journals AMT & GMD
- Financial Support for Authors
- Journal Impact Factor
- Public Relations & Background Information

[Recent Papers](#)

01 | ACP, 06 Feb 2009:
The Cloud Condensation Nuclei (CCN) properties of 2-methyltetrols and C3-C6 polyols from osmolality and surface tension measurements

02 | ACP, 06 Feb 2009:
Airborne measurements of nucleation mode particles II: boreal forest nucleation events

03 | ACP, 06 Feb 2009:
Coupling aerosol-cloud-radiative processes in the WRF-Chem model: Investigating the radiative