

Home

Online Library ACP

- ▣ Recent Final Revised Papers
- ▣ [Volumes and Issues](#)
- ▣ Special Issues
- ▣ Library Search
- ▣ Title and Author Search

Online Library ACPD

Alerts & RSS Feeds

General Information

Submission

Review

Production

Subscription

Comment on a Paper

Impact  
Factor  
4.865

ISI  
indexed



▣ [Volumes and Issues](#) ▣ [Contents of Issue 4](#)

Atmos. Chem. Phys., 4, 923-932, 2004

[www.atmos-chem-phys.net/4/923/2004/](http://www.atmos-chem-phys.net/4/923/2004/)

© Author(s) 2004. This work is licensed under a Creative Commons License.

## On the spectrum of vertically propagating gravity waves generated by a transient heat source

M. J. Alexander<sup>1</sup> and J. R. Holton<sup>2</sup>

<sup>1</sup>Colorado Research Associates Division, NorthWest Research Associates, Inc., Boulder, CO, USA

<sup>2</sup>Department of Atmospheric Sciences, University of Washington, Seattle, WA, USA

**Abstract.** It is commonly believed that cumulus convection preferentially generates gravity waves with tropospheric vertical wavelengths approximately twice the depth of the convective heating. Individual cumulonimbus, however, act as short term transient heat sources (duration 10 to 30min). Gravity waves generated by such sources have broad frequency spectra and a wide range of vertical scales. The high-frequency components tend to have vertical wavelengths much greater than twice the depth of the heating. Such waves have large vertical group velocities, and are only observed for a short duration and at short horizontal distances from the convective source. At longer times and longer distances from the source the dominant wave components have short vertical wavelengths and much slower group velocities, and thus are more likely to be observed even though their contribution to the momentum flux in the upper stratosphere and mesosphere may be less than that of the high frequency waves. These properties of convectively generated waves are illustrated by a linear numerical model for the wave response to a specified transient heat source. The wave characteristics are documented through Fourier and Wavelet analysis, and implications for observing systems are discussed.

▣ [Final Revised Paper](#) (PDF, 1372 KB) ▣ [Discussion Paper](#) (ACPD)

Citation: Alexander, M. J. and Holton, J. R.: On the spectrum of vertically propagating gravity waves generated by a transient heat source, Atmos. Chem. Phys., 4, 923-932, 2004. ▣ [Bibtex](#) ▣ [EndNote](#) [Reference Manager](#)



Search ACP

Library Search

Author Search

News

- ▣ [Sister Journals AMT & GMD](#)
- ▣ [Financial Support for Authors](#)
- ▣ [Journal Impact Factor](#)
- ▣ [Public Relations & Background Information](#)

Recent Papers

01 | ACPD, 27 Feb 2009:  
Effects of boundary layer particle formation on cloud droplet number and changes in cloud albedo from 1850 to 2000

02 | ACPD, 27 Feb 2009:  
A product study of the isoprene+NO<sub>3</sub> reaction

03 | ACPD, 26 Feb 2009:  
Discriminating low frequency components from long range persistent fluctuations in daily atmospheric temperature variability