

Home

Online Library ACP

Recent Final Revised Papers

Volumes and Issues

Special Issues

Library Search

Title and Author Search

Online Library ACPD

Alerts & RSS Feeds

General Information

Submission

Review

Production

Subscription

Comment on a Paper

Impact Factor  
4.865

ISI  
indexed



Volumes and Issues Contents of Issue 1

Atmos. Chem. Phys., 8, 129-139, 2008

www.atmos-chem-phys.net/8/129/2008/

© Author(s) 2008. This work is licensed under a Creative Commons License.

## Formation and characteristics of ions and charged aerosol particles in a native Australian Eucalypt forest

T. Suni<sup>1</sup>, M. Kulmala<sup>1</sup>, A. Hirsikko<sup>1</sup>, T. Bergman<sup>2</sup>, L. Laakso<sup>1</sup>, P. P. Aalto<sup>1</sup>, R. Leuning<sup>3</sup>, H. Cleugh<sup>3</sup>, S. Zegelin<sup>3</sup>, D. Hughes<sup>3</sup>, E. van Gorsel<sup>3</sup>, M. Kitchen<sup>3</sup>, M. Vana<sup>4</sup>, U. Hörrak<sup>4</sup>, S. Mirme<sup>4</sup>, A. Mirme<sup>4</sup>, S. Sevanto<sup>1</sup>, J. Twining<sup>5</sup>, and C. Tardos<sup>5</sup>

<sup>1</sup>Department of Physical Sciences, P.O.Box 64, FIN-00014 University of Helsinki, Finland

<sup>2</sup>Center for High-Performance Computing and Networking CSC, PO. BOX 405, 02101 Espoo, Finland

<sup>3</sup>CSIRO Marine and Atmospheric Research, GPO Box 1666, Canberra ACT 2601, Australia

<sup>4</sup>Institute of Physics, University of Tartu, Ülikooli Str 18., EE2400 Tartu, Estonia

<sup>5</sup>Institute for Environmental Research, ANSTO, PMB 1, Menai NSW 2234, Australia

**Abstract.** Biogenic aerosol formation is likely to contribute significantly to the global aerosol load. In recent years, new-particle formation has been observed in various ecosystems around the world but hardly any measurements have taken place in the terrestrial Southern Hemisphere. Here, we report the first results of atmospheric ion and charged particle concentrations as well as of new-particle formation in a Eucalypt forest in Tumbarumba, South-East Australia, from July 2005 to October 2006. The measurements were carried out with an Air Ion Spectrometer (AIS) with a size range from 0.34 to 40 nm. The Eucalypt forest was a very strong source of new aerosol particles. Daytime aerosol formation took place on 52% of days with acceptable data, which is 2–3 times as often as in the Nordic boreal zone. Average growth rates for negative/positive 1.5–3 nm particles during these formation events were 2.89/2.68 nmh<sup>-1</sup>, respectively; for 3–7 nm particles 4.26/4.03, and for 7–20 nm particles 8.90/7.58 nmh<sup>-1</sup>, respectively. The growth rates for large ions were highest when the air was coming from the native forest which suggests that the Eucalypts were a strong source of condensable vapours. Average concentrations of cluster ions (0.34–1.8 nm) were 2400/1700 cm<sup>-3</sup> for negative/positive ions, very high compared to most other measurements around the world. One reason behind these high concentrations could be the strong radon efflux from the soils around the Tumbarumba field site. Furthermore, comparison between night-time and daytime concentrations supported the view that cluster ions are produced close to the surface within the boundary layer also at night but that large ions are mostly produced in daytime. Finally, a previously unreported phenomenon, nocturnal aerosol formation, appeared in 32% of the analysed nights but was clustered almost entirely within six months from summer to autumn in 2006. From January to May, nocturnal formation was 2.5 times as frequent as daytime formation. Therefore, it appears that in summer and autumn, nocturnal production was the major mechanism for aerosol formation in Tumbarumba.

Search ACP

Library Search

Author Search

News

- Sister Journals AMT & GMD
- Financial Support for Authors
- Journal Impact Factor
- Public Relations & Background Information

Recent Papers

01 | ACPD, 03 Nov 2008:  
Technical Note: A new method for the Lagrangian tracking of pollution plumes from source to receptor using gridded model output

02 | ACPD, 03 Nov 2008:  
Characterisation of episodic aerosol types over the Australian continent

03 | ACPD, 03 Nov 2008:  
Evidence of mineral dust altering cloud microphysics and precipitation

04 | ACP, 03 Nov 2008:

▣ [Final Revised Paper](#) (PDF, 548 KB) ▣ [Discussion Paper](#) (ACPD)

Citation: Suni, T., Kulmala, M., Hirsikko, A., Bergman, T., Laakso, L., Aalto, P. P., Leuning, R., Cleugh, H., Zegelin, S., Hughes, D., van Gorsel, E., Kitchen, M., Vana, M., Hörrak, U., Mirme, S., Mirme, A., Sevanto, S., Twining, J., and Tardos, C.: Formation and characteristics of ions and charged aerosol particles in a native Australian Eucalypt forest, Atmos. Chem. Phys., 8, 129-139, 2008. ▣ [Bibtex](#) ▣ [EndNote](#) [Reference Manager](#)