

Home

Online Library ACP

- Recent Final Revised Papers
- [Volumes and Issues](#)
- Special Issues
- Library Search
- Title and Author Search

Online Library ACPD

Alerts & RSS Feeds

General Information

Submission

Review

Production

Subscription

Comment on a Paper

Impact
Factor
4.865

ISI
indexed



[Volumes and Issues](#) [Contents of Issue 11](#)

Atmos. Chem. Phys., 5, 3139-3151, 2005
www.atmos-chem-phys.net/5/3139/2005/

© Author(s) 2005. This work is licensed
under a Creative Commons License.

Impact of mixing and chemical change on ozone-tracer relations in the polar vortex

R. Müller¹, S. Tilmes^{1,*}, P. Konopka¹, J.-U. Grooß², and H.-J. Jost

¹ICG I, Forschungszentrum Jülich, 52425 Jülich, Germany

²Bay Area Environmental Research Institute, Sonoma, CA, USA

*now at: ACD, NCAR, Boulder, CO, USA

Abstract. Tracer-tracer relations have been used for a long time to separate physico-chemical change from change caused by transport processes. In particular, for more than a decade, ozone-tracer relations have been used to quantify chemical ozone loss in the polar vortex. The application of ozone-tracer relations for quantifying ozone loss relies on two hypotheses: that a compact ozone-tracer relation is established in the 'early' polar vortex and that any change of the ozone-tracer relation in the vortex over the course of winter is caused predominantly by chemical ozone loss. Here, we revisit this issue by analysing various sets of measurements and the results from several models. We find that mixing across the polar vortex edge impacts ozone-tracer relations in a way that may solely lead to an 'underestimation' of chemical ozone loss and not to an overestimation. Further, differential descent in the vortex and internal mixing has only a negligible impact on ozone loss estimates. Moreover, the representation of mixing in three-dimensional atmospheric models can have a substantial impact on the development of tracer relations in the model. Rather compact ozone-tracer relations develop – in agreement with observations – in the vortex of a Lagrangian model (CLaMS) where mixing is anisotropic and driven by the deformation of the flow. We conclude that, if a reliable 'early vortex' reference can be obtained and if vortex measurements are separated from mid-latitude measurements, ozone-tracer relations constitute a reliable tool for the quantitative determination of chemical ozone loss in the polar vortex.

[Final Revised Paper](#) (PDF, 3630 KB)

Citation: Müller, R., Tilmes, S., Konopka, P., Grooß, J.-U., and Jost, H.-J.: Impact of mixing and chemical change on ozone-tracer relations in the polar vortex, Atmos. Chem. Phys., 5, 3139-3151, 2005. [Bibtex](#) [EndNote](#) [Reference Manager](#)

Search ACP

Library Search

Author Search

News

- [Sister Journals AMT & GMD](#)
- [Financial Support for Authors](#)
- [Journal Impact Factor](#)
- [Public Relations & Background Information](#)

Recent Papers

01 | ACPD, 14 Jan 2009:
The impact of tropical
recirculation on polar
composition

02 | ACPD, 14 Jan 2009:
Lightning NO_x emissions over
the USA investigated using
TES, NLDN, LRLDN, IONS data
and the GEOS-Chem model

03 | ACP, 14 Jan 2009:
Properties of aerosols and
their wet deposition in the
arctic spring during
ASTAR2004 at Ny-Alesund,
Svalbard