

地球动力学★地震学★地磁学

利用1996年丽江地震序列反演震区应力状态

张永庆^{1,2}, 谢富仁¹, Susanna J. Gross³

1 中国地震局地壳应力研究所, 北京 100085

2 中国地震局地质研究所, 北京 100029

3 Cooperative Institute for Research in the Environmental Sciences, University of Colorado, Boulder, Colorado 80309, USA

收稿日期 2008-5-20 修回日期 2009-3-24 网络版发布日期 2009-4-20 接受日期

摘要 本文以1996年丽江 M_S 7.0地震为例, 以地震目录和强震震源模型为输入参数, 将力学计算与统计分析相结合, 反演计算了1986~1996年间丽江震区 $2.5^\circ \times 3.0^\circ$ 空间范围内的地壳应力状态. 结果表明, 丽江地区应力场最大主应力为近南北向, 方位角为 355° , 最小主应力方位角为 241° , 二者倾角为近水平, 应力形因子为0.398, 最大主应力大小约为53 MPa, 远小于完整岩石的强度. 该方法可获得地壳深部绝对应力大小、有效摩擦系数, 反演获得的震区地壳深部应力状态的方向性特征与震源机制解结果基本一致; 该研究表明地震活动的空间分布可以为我们提供有关强震孕震的地壳深部应力状态信息. 这些结果对了解震区应力场具有理论上和实践上的意义.

关键词 [丽江地震](#) [地震序列](#) [反演](#) [震区](#) [构造应力状态](#)

分类号 [P541](#)

DOI: [10.3969/j.issn.0001-5733.2009.04.019](#)

Background stress state estimated from 1996 Lijiang earthquake sequence

ZHANG Yong-Qing^{1,2}, XIE Fu-Ren¹, Susanna J. Gross³

1 Institute of Crustal Dynamics, China Earthquake Administration, Beijing 100085, China

2 Institute of Geology, China Earthquake Administration, Beijing 100029, China

3 Cooperative Institute for Research in the Environmental Sciences, University of Colorado, Boulder, Colorado 80309, USA

Received 2008-5-20 Revised 2009-3-24 Online 2009-4-20 Accepted

Abstract Estimates of the tectonic stress state including direction and magnitude of principal stress are derived from the regional seismicity before and after 1996, M_S 7.0 Lijiang earthquake. This technique was originally applied to the Landers aftershock sequence (Gross and Kisslinger, 1997). Two kinds of input include a combined catalogue of events before and after the Lijiang earthquake, and a source model derived from the inversion of body wave. The stress field in source region was calculated using a 3-D dislocation model, and the best stress state is statistically determined by evaluating the relativity between stress change and seismicity change. The result shows that the azimuth of maximum principal stress in 10 km depth is 355° , and the magnitude of maximum principal stress is 53MPa, which is much smaller than the value estimated from the intact rock at that depth. Inversions for the best fitting background stress state are consistent with focal mechanism solution (with a 12° degree difference). This study shows that the spatial distribution of seismicity can present stress state information of source area. Although the estimations of the magnitude of background stress, gradient of stress with depth and effective coefficient of friction are quite uncertain, and need further research to confirm, the result provides the dynamic information of stress loading on the faults, which is useful for the further research on stress environment of seismogenic area.

Key words [Lijiang earthquake](#); [Earthquake sequence](#); [Inversion](#); [Source area](#); [Tectonic stress state](#)

通讯作者:

张永庆 whyzyq@yahoo.com.cn

作者个人主页: 张永庆^{1,2}; 谢富仁¹; Susanna J. Gross³

扩展功能

本文信息

▶ [Supporting info](#)

▶ [PDF](#) (1785KB)

▶ [\[HTML全文\]](#) (0KB)

▶ [参考文献](#)

服务与反馈

▶ [把本文推荐给朋友](#)

▶ [加入我的书架](#)

▶ [加入引用管理器](#)

▶ [引用本文](#)

▶ [Email Alert](#)

▶ [文章反馈](#)

▶ [浏览反馈信息](#)

相关信息

▶ [本刊中 包含“丽江地震”的 相关文章](#)

▶ 本文作者相关文章

• [张永庆](#)

•

• [谢富仁](#)

• [Susanna J Gross](#)