

Home**Online Library HESS**

- Recent Final Revised Papers
- [Volumes and Issues](#)
- Special Issues
- Library Search
- Title and Author Search

Online Library HESSD**Alerts & RSS Feeds****General Information****Submission****Review****Production****Subscription****Comment on a Paper**

ARCHIVED IN

**Volumes and Issues** ■ **Contents of Issue 3**

Hydrol. Earth Syst. Sci., 6, 363-374, 2002

www.hydrol-earth-syst-sci.net/6/363/2002/

© Author(s) 2002. This work is licensed under a Creative Commons License.

Spatial variability of throughfall water and chemistry and forest floor water content in a Douglas fir forest stand

K. J. Raat¹, G. P. J. Draaijers², M. G. Schaap³, A. Tietema¹, and J. M. Verstraten¹¹Centre for Geo-Ecological Research (ICG), Institute for Biodiversity and Ecosystem Dynamics (IBED) - Physical Geography, Universiteit van Amsterdam, Nieuwe Achtergracht 166, NL-1018WV Amsterdam, The Netherlands²TNO Institute of Environmental Sciences, Energy Research and Process Innovation (TNO-MEP), P.O. Box 342, NL-7300 AH Apeldoorn, The Netherlands³George E. Brown Jr. Salinity Laboratory, USDA-ARS, 450 W. Big Springs Road, Riverside, CA 92507, USA

Email for corresponding author: k.raat@science.uva.nl

Abstract. This study focuses on spatial variability of throughfall water and chemistry and forest floor water content within a Douglas fir (*Pseudotsuga menziesii*, Franco L.) forest plot. Spatial patterns of water and chemistry (NH_4^+ , NO_3^- , SO_4^{2-} , Cl^- , Mg^{2+} , Ca^{2+} , Na^+ and K^+) were compared and tested for stability over time. The spatial coefficient of variation (CV) was between 18 and 26% for amounts of throughfall water and ions, and 17% for forest floor water content. Concentrations and amounts of all ions were correlated significantly. Ion concentrations were negatively correlated with throughfall water amounts, but, except for NH_4^+ , there was no such relation between throughfall water and ion amounts. Spatial patterns of throughfall water fluxes and forest floor water contents were consistent over time; patterns of ion fluxes were somewhat less stable. Because of the spatial variability of forest floor thickness and drainage, it was not possible to relate patterns in throughfall water directly to patterns in water content. The spatial variability of throughfall nitrogen and forest floor water contents can cause significant variability in NO_3^- production within the plot studied.

Keywords: nutrient throughfall, forest floor water, spatial variability, time-stability, nitrogen

■ [Final Revised Paper](#) (PDF, 736 KB)

Citation: Raat, K. J., Draaijers, G. P. J., Schaap, M. G., Tietema, A., and Verstraten, J. M.: Spatial variability of throughfall water and chemistry and forest floor water content in a Douglas fir forest stand, Hydrol. Earth Syst. Sci., 6, 363-374, 2002. ■ [Bibtex](#) ■ [EndNote](#) ■ [Reference Manager](#)

**Search HESS** Library Search Author Search **News**

- New Service Charges
- Financial Support for Authors
- ISI Impact Factor: 2.270

Recent Papers

01 | HESSD, 17 Mar 2009:
A general real-time formulation for multi-rate mass transfer problems

02 | HESSD, 16 Mar 2009:
Calibration of a crop model to irrigated water use using a genetic algorithm

03 | HESSD, 16 Mar 2009:
A Bayesian approach to estimate sensible and latent heat over vegetation

04 | HESS, 13 Mar 2009:
Soil moisture retrieval through a merging of multi-