

Home

Online Library HESS

- ▣ Recent Final Revised Papers
- ▣ [Volumes and Issues](#)
- ▣ Special Issues
- ▣ Library Search
- ▣ Title and Author Search

Online Library HESSD

Alerts & RSS Feeds

General Information

Submission

Review

Production

Subscription

Comment on a Paper



▣ [Volumes and Issues](#) ▣ [Contents of Issue 1](#)

Hydrol. Earth Syst. Sci., 12, 159-175, 2008  
[www.hydrol-earth-syst-sci.net/12/159/2008/](http://www.hydrol-earth-syst-sci.net/12/159/2008/)

© Author(s) 2008. This work is licensed under a Creative Commons License.

## Strong increases in flood frequency and discharge of the River Meuse over the late Holocene: impacts of long-term anthropogenic land use change and climate variability

P. J. Ward<sup>1</sup>, H. Renssen<sup>1</sup>, J. C. J. H. Aerts<sup>2</sup>, R. T. van Balen<sup>1</sup>, and J. Vandenberghe<sup>1</sup>

<sup>1</sup>Institute of Earth Sciences, Faculty of Earth and Life Sciences, VU University Amsterdam, De Boelelaan 1085, 1081 HV Amsterdam, The Netherlands

<sup>2</sup>Institute for Environmental Studies, Faculty of Earth and Life Sciences, VU University Amsterdam, De Boelelaan 1085, 1081 HV Amsterdam, The Netherlands

**Abstract.** In recent years the frequency of high-flow events on the Meuse (northwest Europe) has been relatively great, and flooding has become a major research theme. To date, research has focused on observed discharge records of the last century and simulations of the coming century. However, it is difficult to delineate changes caused by human activities (land use change and greenhouse gas emissions) and natural fluctuations on these timescales. To address this problem we coupled a climate model (ECBilt-CLIO-VECODE) and a hydrological model (STREAM) to simulate daily Meuse discharge in two time-slices: 4000–3000 BP (natural situation), and 1000–2000 AD (includes anthropogenic influence). For 4000–3000 BP the basin is assumed to be almost fully forested; for 1000–2000 AD we reconstructed land use based on historical sources. For 1000–2000 AD the simulated mean annual discharge ( $260.9 \text{ m}^3 \text{ s}^{-1}$ ) is significantly higher than for 4000–3000 BP ( $244.8 \text{ m}^3 \text{ s}^{-1}$ ), and the frequency of large high-flow events (discharge  $>3000 \text{ m}^3 \text{ s}^{-1}$ ) is higher (recurrence time decreases from 77 to 65 years). On a millennial timescale almost all of this increase can be ascribed to land use changes (especially deforestation); the effects of climatic change are insignificant. For the 20th Century, the simulated mean discharge ( $270.0 \text{ m}^3 \text{ s}^{-1}$ ) is higher than in any other century studied, and is ca. 2.5% higher than in the 19th Century (despite an increase in evapotranspiration). Furthermore, the recurrence time of large high-flow events is almost twice as short as under natural conditions (recurrence time decreases from 77 to 40 years). On this timescale climate change (strong increase in annual and winter precipitation) overwhelmed land use change as the dominant forcing mechanism.

▣ [Final Revised Paper](#) (PDF, 3806 KB) ▣ [Discussion Paper](#) (HESSD)

Citation: Ward, P. J., Renssen, H., Aerts, J. C. J. H., van Balen, R. T., and Vandenberghe, J.: Strong increases in flood frequency and discharge of the River Meuse over the late Holocene: impacts of long-term anthropogenic land use change and climate variability, Hydrol. Earth Syst. Sci., 12, 159-175, 2008. ▣ [Bibtex](#) ▣ [EndNote](#) ▣ [Reference Manager](#)



Search HESS

Library Search

Author Search

News

- ▣ [New Service Charges](#)
- ▣ [Financial Support for Authors](#)
- ▣ [ISI Impact Factor: 2.270](#)

Recent Papers

01 | HESSD, 28 Apr 2009: Integrating field and numerical modeling methods for applied urban karst hydrogeology

02 | HESSD, 28 Apr 2009: Analyzing the relationship between peak runoff discharge and land-use pattern – a spatial optimization approach

03 | HESSD, 27 Apr 2009: Dynamically vs. empirically downscaled medium-range precipitation forecasts

