



## Full Text View

[Volume 29, Issue 8 \(August 1999\)](#)

### Journal of Physical Oceanography

Article: pp. 2113–2116 | [PDF \(68K\)](#)

## Reply

**Ray Q. Lin**

*Hydromechanics Directorate, David Taylor Model Basin, Carderock Division, West Bethesda, Maryland*

**Will Perrie**

*Ocean Sciences Division, Fisheries and Oceans Canada, Bedford Institute of Oceanography, Dartmouth, Nova Scotia, Canada*

(Manuscript received February 11, 1999)

DOI: 10.1175/1520-0485(1999)029<2113:R>2.0.CO;2

### ABSTRACT

#### 1. Introduction

In the accompanying comment, Kalmykov expresses two concerns regarding the work of [Lin and Perrie \(hereafter LP\)](#) 1) that the numerical simulations of five-wave interactions by [Kalmykov](#) were not cited by LP and 2) that, because the results of Kalmykov are qualitatively different from those of LP, therefore the conclusions of LP cannot be justified.

The authors apologize for not citing the work of [Kalmykov](#). These papers constitute important early attempts to compute the five-wave interactions. Unfortunately, the later [Kalmykov](#) paper did not become available until after the LP paper had gone to press.

The authors' major concern in this reply is to demonstrate that the conclusions of LP are *correct*. The five-wave interactions, as described by LP, are for finite amplitude waves with a narrow spectral spreading and large peakedness  $\gamma$ , rather than small amplitude waves with a broad spectral spreading and small  $\gamma = 1$ , which is the approach taken by Kalmykov's papers.

#### 2. Scientific background

First, the dominance of the five-wave interactions direct or inverse cascades depends on the shape of the spectrum. Five-wave interactions can cause both energy transfer from low to high frequencies (the direct cascade) and energy transfer from high frequencies to low frequencies (the inverse cascade). These cascades are

#### Table of Contents:

- [Introduction](#)
- [Scientific background](#)
- [Numerical simulations](#)
- [Previous work](#)
- [Summary](#)
- [REFERENCES](#)
- [FIGURES](#)

#### Options:

- [Create Reference](#)
- [Email this Article](#)
- [Add to MyArchive](#)
- [Search AMS Glossary](#)

#### Search CrossRef for:

- [Articles Citing This Article](#)

#### Search Google Scholar for:

- [Ray Q. Lin](#)
- [Will Perrie](#)

shown in [Fig. 1](#). Following LP, a dominant inverse cascade is shown in [Fig. 1a](#), assuming a narrow (Hasselmann–Mitsuyasu) spectral spreading, as motivated by [Hasselmann et al. \(1980\)](#) from JONSWAP analysis,

$$D(\theta, f) = I(s) \cos^{2s}[(\theta - \theta_{\max})/2], \quad (1)$$

where  $I(s)$  is a normalization factor and  $s$  is parameterized as

$$s = \begin{cases} 6.97(f/f_p)^{4.06} & \text{for } f/f_p < 1.05. \\ 9.77(f/f_p)^\mu & \text{for } f/f_p \geq 1.05. \end{cases} \quad (2)$$

Here,  $\mu$  is a weakly dependent function of wave age,  $U_{10}/C_p$ , which satisfies the relation

$$\mu = 2.33 - 1.45 \left( \frac{U_{10}}{C_p} - 1.17 \right),$$

where  $C_p$  is the phase speed at  $f_p$ . A similar form was proposed by [Mitsuyasu et al. \(1975\)](#).

However, if the spectrum is very broad in angle, for example  $\cos^2\theta$ , then the dominance of a direct cascade, as suggested by Kalmykov's comment and papers, is correct. In [Fig. 1b](#), we give five-wave interactions direct and inverse cascades for the latter spreading, showing the dominance of the direct cascade. However, in this case, assuming deep water waves that are not steep, LP and Kalmykov's papers all suggest that four-wave interactions should dominate over five-wave interactions, implying that inverse cascades due to four-wave interactions would be a dominant feature.

Second, four-wave interactions are local interactions, which are proportional to  $d^3$ , where  $d$  is the width of the spectrum, whereas five-wave interactions are global interactions that are not significantly affected by  $d$ , as presented in LP. Therefore, the five-wave interactions allow the study of finite-amplitude narrow spectral interactions, which may be able to obtain significant three-dimensional patterns, as described by [McLean's \(1982a,b\)](#) instability analysis, [Su's \(1982a,b\)](#) experimental analysis, [Long et al.'s \(1994\)](#) observations, and the long-lived patterns of [Shrira et al. \(1996\)](#).

Third, as discussed in [Zakharov \(1991\)](#), the direct cascades cause the energy transfer from low to high frequency, whereas the inverse cascades cause the energy transfer from high to low frequency. The latter can cause waves to grow and the peak frequency to downshift to lower frequencies, whereas the former causes waves to break. Therefore, inverse cascades, as given by the finite-amplitude, five-wave interactions of LP, are central to the long timescale, three-dimensional mechanics, as observed by [Su \(1982a,b\)](#), [McLean \(1982a,b\)](#), [Long et al. \(1994\)](#), and [Shrira et al. \(1996\)](#), whereas associated direct cascades are central to the short timescale mechanics for remote sensing and radar scattering. As four-wave interactions are primarily two-dimensional, whereas five-wave interactions are primarily three-dimensional, the four-wave interactions, by themselves, cannot provide a comprehensive description of these three-dimensional patterns of [Su \(1982a,b\)](#). By contrast, the five-wave interactions of [Kalmykov \(1999, p. 2\)](#) are only significant in shallow water. Unless waves are extremely steep, they are only associated with direct cascades and they are always much smaller than the corresponding four-wave interactions. Therefore, the five-wave interactions of the comment inhibit wave growth because of their dominant direct cascades, and they cannot account for the finite-amplitude three-dimensional patterns of [Su \(1982a,b\)](#).

Fourth, recently [Lin and Chubb \(1999\)](#) found significant direct cascades by weakly nonlinear four-wave interactions when the initial spectrums are split so the observed direct cascades at the tail of the spectrum may be due to four-wave interactions instead of five-wave interactions.

Finally, it is well known that five-wave interactions can be significant only when the wave steepness is greater than about 0.3, as given by [Su \(1982a,b\)](#), [McLean \(1982a,b\)](#), [Shrira et al. \(1996\)](#), and LP. At the bottom of p. 2, [Kalmykov \(1999\)](#) states that "This instability II (3D) was triggered by instability I (2D) or Benjamin-Feir instability." This is inconsistent with the comment (i) at the top of p. 2, where Kalmykov suggests that these four-wave resonant conditions cannot be satisfied in shallow water and that five-wave interactions would result, and (ii) on p. 6 of the comment, where Kalmykov uses scaling arguments to suggest that even for very steep waves in very shallow water, five-wave interactions should be much less important than four-wave interactions. Moreover, we note that the mechanics by which four- and five-wave interactions are coupled to generate 3D wave-wave interactions patterns was suggested by [Su \(1982a,b\)](#) for deep water waves, not for shallow water waves.

### 3. Numerical simulations

We have some concerns regarding [Kalmykov's \(1993, 1995, 1997, 1998\)](#) results.

1. A major problem is that in the first three of these papers, the nonlinear transfer due to five-wave interactions is strongly *asymmetric*. Figure 3 of [Kalmykov \(1997\)](#) notes that the energy is moved from the spectral maximum  $\theta_o = 0$  to lateral waves at  $\theta_2 = 60^\circ$  and  $\theta_1 = -90^\circ$ . By contrast [Kalmykov \(1998\)](#) gives *symmetric* lateral transfer to waves at  $\theta = \pm 70^\circ$ , which is similar to the lateral transfer angle presented by LP. No discussion is given in regard to this lateral transfer. Thus, the results of [Kalmykov \(1998\)](#) appear inconsistent with the results of [Kalmykov \(1993, 1995, 1997\)](#).
2. It is difficult to quantitatively compare the magnitudes of Kalmykov's results because of the nondimensionalization he has implemented. Even in nondimensional units, his results show some variation in his four papers cited above. For example, in dimensionless variables, five-wave interactions have a maximal magnitude of about 0.05 in [Kalmykov \(1995, 1997\)](#), which becomes about 18 in [Kalmykov \(1999\)](#) and [Kalmykov \(1998\)](#). [Kalmykov \(1998\)](#) does indicate the factor to convert these values to dimensional values, which then gives maximal magnitudes of order  $10^{-5}$ , similar to LP maximal values. However, a detailed direct comparison between LP and Kalmykov's results is difficult to make because Kalmykov uses a Pierson–Moskowitz input spectrum, which corresponds to very old sea state conditions, whereas LP use JONSWAP input spectrum with peak frequency at 0.3 Hz, corresponding to actively growing, evolving, young sea state conditions. Moreover, Kalmykov's directional spreading is  $\cos^2\theta$  for  $|\theta| \leq \pi/2$ , which is significantly broader than the Hasselmann–Mitsuyasu spreading in LP, as described above. Although [Kalmykov \(1999\)](#) claims to use a JONSWAP input spectrum, it is, in fact, the old sea state spectrum (Pierson–Moskowitz) of Fig. 2a in [Kalmykov \(1998\)](#).
3. [Young et al. \(1996\)](#) suggest that shallow water spectra are more broad than deep water spectra, based on an extensive data analysis, comparing his shallow lake data to [Donelan et al.'s \(1985\)](#) deep water data from Lake Ontario. Using a broadly spreading spectrum, for example  $\cos^2\theta$ , [Kalmykov \(1998\)](#) suggests that the effect of shallower water is to *narrow* the spectrum, which is therefore contradictory to [Young et al.'s \(1996\)](#) observations.
4. Numerical simulation of the nonlinear energy transfer depends on both the transfer coefficient and the integration method. As Kalmykov claims his transfer coefficient is taken from [Krasitski \(1993\)](#), his simulation should be mostly correct. However, problems occur in his integration method and related assumptions. [Kalmykov \(1997\)](#) assumes the five-wave resonance conditions are given by  $3k_o = k_4 + k_5$  and  $3\omega_o = \omega_4 + \omega_5$ , where  $k_o = \omega_o = 1$ , which is impossible for five-wave interactions. Unlike four-wave interactions, five-wave interactions are global interactions, which means that the five interacting waves do not need to have comparable wavelengths. This assumption makes [Kalmykov's \(1997\)](#) solutions qualitatively different from the real solutions.

#### 4. Previous work

[Kalmykov \(1999\)](#) claims that all previous works, such as [Dyachenko et al. \(1994\)](#), [Krasitski \(1993, 1994\)](#), [Meiron et al. \(1982\)](#), [Shrira et al. \(1996\)](#), [McLean \(1982a,b\)](#), [Stiassnie and Shemer \(1984, 1987\)](#), and [Su \(1982a,b\)](#) support his results, including his claim that five-wave interactions support only direct cascades. Unfortunately, this is not correct. [Dyachenko et al. \(1994\)](#), [Krasitski \(1993, 1994\)](#), [Meiron et al. \(1982\)](#), and [Stiassnie and Shemer \(1984, 1987\)](#) basically studied the nonlinear transfer coefficient, not the numerical simulation. They concluded that five-wave interactions will generate three-dimensional wave–wave interactions. They did not discuss the ability of five-wave interactions to support inverse cascades or direct cascades.

[Shrira et al. \(1996\)](#) suggested that class II instability (five-wave interactions, seven-wave interactions, and so on) generate horseshoe patterns. However, inverse cascades and direct cascades are both present in horseshoe patterns, and [Shrira et al. \(1996\)](#) did not suggest that only direct cascades are present, as [Kalmykov \(1999\)](#) claims. Moreover, [Shrira et al. \(1996\)](#) did suggest that wave steepness needs to be greater than 0.33 in order for the generation of the three-dimensional patterns, which is approximately the same as LP's criterion for the dominance of five-wave interactions. Thus, five-wave interactions must support inverse cascades, as suggested by LP. Otherwise, if five-wave interactions only support direct cascades and if they dominate over four-wave interactions, as found by LP, then three-dimensional wave patterns will not be observable because it is impossible for waves to grow and evolve.

[McLean \(1982a,b\)](#) used a global linear method to study five-wave instability. He found that five-wave interactions can be greater than four-wave interactions when wave steepness is greater than 0.28. [Su \(1982a,b\)](#) first observed the three-dimensional wave–wave interactions in his data. These results agree with the finite-amplitude analysis of LP and disagree with [Kalmykov \(1993, 1995, 1997, 1998\)](#). [Su and Green's \(1984\)](#) analysis, based on their experimental results, suggests that the two-dimensional instabilities may trigger the three-dimensional instabilities in deep water. However, this approach does not apply to shallow water, because four-wave interactions become rapidly less important as water depth decreases, as shown in [Lin and Perrie \(1997b\)](#).

Recently, [Lin and Su \(1999\)](#) showed that a coupling of four- and five-wave interactions can cause three-dimensional wave-wave interactions in deep water, when the spectrum is narrow and wave steepness is significant. The results agree well with LP. When inverse cascades occur, transferring energy from high to low frequencies, then waves can only grow and the coupling of four- and five-wave interactions can occur.

The analysis of [Long et al. \(1994\)](#) considers three sets of observed wave spectra in shallow water. The dominant directions of the wave spectra distributions are about  $60^\circ$  from the downwind direction. The wave amplitudes are about 4 m high. These are finite-amplitude waves. While LP can provide a good explanation for these wave spectra phenomena, because their transfer to lateral waves by the five-wave interactions is at  $\pm 60^\circ$ , other mechanisms such as the Phillips mechanism for wave generation or the five-wave interactions of [Kalmykov \(1993, 1995, 1997, 1998\)](#), cannot explain these phenomena.

[Kalmykov \(1999\)](#) states, “For any odd number of the wave interactions, the energy transfer is a direct cascade: from low frequency to high, while for any even number we get an inverse cascade ([Zakharov 1998](#)).” We could not find this stated or implied in [Zakharov \(1998\)](#). Both direct and inverse cascades are computed in [Resio and Perrie \(1991\)](#) for four-wave interactions, in accord with the theoretical study by [Zakharov \(1991\)](#). Moreover, Prof. Zakharov has discussed the nonlinear wave-wave interactions of LP several times with us: he has never suggested that odd (even) number wave-wave interactions should have only a direct (inverse) cascade.

## 5. Summary

We have shown that five-wave interactions should include both direct and indirect cascades, depending on the spectral spreading. This follows from the existence of the finite-amplitude long-lived, three-dimensional patterns, as observed by [Su \(1982a,b\)](#), and the associated analysis of [McLean \(1982a,b\)](#). Therefore, following [Zakharov \(1991\)](#), these features should be generated by inverse cascades, not direct cascades, as presented by LP. In this sense, we have shown that Kalmykov’s results contradict the instability analysis of [McLean \(1982a,b\)](#) and the observational data of [Su \(1982a,b\)](#).

[Kalmykov’s \(1993, 1995, 1997\)](#) earlier results are strongly asymmetrical whereas his most recent results, shown in [Kalmykov \(1998\)](#), are symmetrical. The former are unphysical, because there is no obvious reason for asymmetry. Kalmykov gives no apparent discussion of this asymmetry/symmetry.

When the spectrum is broad, such as  $\cos^m\theta$ , and when peakedness  $\gamma$  is small, the direct cascade tends to be more dominant, as suggested by [Kalmykov \(1998\)](#) and presented in [Fig. 1b](#). When the spectrum is narrower, as parameterized by the Hasselmann–Mitsuasyu spreading and when  $\gamma$  is large, the inverse cascade tends to dominate, as shown in [Fig. 1a](#) and in LP.

## Acknowledgments

This work has been supported by the Office of Naval Research under the Wave Prediction Base Enhancement Programs. W. Perrie was supported by the Canadian Federal Panel on Energy Research and Development under Project 534201.

---

## REFERENCES


- Donelan, M. A., J. Hamilton, and W. H. Hui, 1985: Directional spectra of wind-generated waves. *Philos. Trans. Roy. Soc. London A*, **315**, 509–562.
- Dyachenko, A. I., Y. V. Lvov, and V. E., Zakharov, 1994: Five-wave interaction on the surface of deep fluid. *Physica D*, **87**, 233–261.
- Hasselmann, D. E., M. Dunkel, and J. A. Ewing, 1980: Directional wave spectral observed during JONSWAP 1973. *J. Phys. Oceanogr.*, **10**, 1264–1280. [Find this article online](#)
- Kalmykov, V. A., 1993: Numerical calculation of the nonlinear transfer of energy in the spectra of surface gravity waves by 5 wave resonant interactions. *Dokl. Akad. Nauk Ukraini*, **8**, 101–104.
- , 1995: Estimates of energy transfer in the spectra of surface gravity waves by nonlinearity of high order. *Dokl. Akad. Nauk Ukraini*, **11**, 87–89.
- , 1997: Numerical calculation of nonlinear transfer of energy in spectra of surface gravity waves from the five wave resonant interactions. *The Air-Sea Interface Radio and Acoustic Sensing, Turbulence and Wave Dynamics*, M. A. Donelan, W. H. Hui, and W. J.

- ,1998: Renergy transfer in the spectrum of surface gravity waves by resonance five wave-wave interactions. *Amer. Math. Soc. Transl.*, **182**, 83–94..
- ,1999: Comments on “A new coastal wave model. Part V: Five-wave interactions.” *J. Phys. Oceanogr.*, **29**, 2110–2112.. [Find this article online](#)
- Krasitski, V. P., 1993: 5 wave kinetic equation for surface gravity waves. *Marine Hydrophys. J.*, **6**, 17–25..
- ,1994: On reduced equations in the Hamiltonian theory of weakly nonlinear surface waves. *J. Fluid Mech.*, **272**, 1–20..
- Lin, R.-Q., and W. Perrie, 1997a: A new coastal wave model. Part V: Five-wave interactions. *J. Phys. Oceanogr.*, **27**, 2169–2186.. [Find this article online](#)
- ,and — ,1997b: *On the Mathematics and Approximation of the Nonlinear Wave-Wave Interactions*. *Advances in Fluid Mechanics*, Computational Mechanics Publications, Vol. 16, 27 pp..
- ,and M.-Y. Su, 1999: Three-dimensional wave-wave interactions by coupling 4-wave and 5-wave interactions in deep water. *J. Proc. Int. Offshore and Polar Engineering Conf.*, in press..
- Long, S. R., N. E. Huang, M. Mollo-Christensen, F. C. Jackson, and G. L. Greenaert, 1994: Directional wind wave development. *Geophys. Res. Lett.*, **12**, 2503–2506..
- McLean, J. W., 1982a: Instabilities of finite amplitude water waves. *J. Fluid Mech.*, **114**, 315–330..
- ,1982b: Instabilities of finite amplitude gravity waves on water of finite depth. *J. Fluid Mech.*, **114**, 331–341..
- Meiron, D. I., P. G. Saffman, and H. C. Yuen, 1982: Calculation of the steady three-dimensional deep water waves. *J. Fluid Mech.*, **124**, 109–121..
- Mitsuyasu, H., F. Tasai, T. Suhara, S. Mizuno, M. Ohkuso, T. Honda, and K. Rikiishi, 1975: Observations of the directional spectra of ocean waves using a cloverleaf buoy. *J. Phys. Oceanogr.*, **5**, 750–760.. [Find this article online](#)
- Resio, D., and W. Perrie, 1991: A numerical study of nonlinear energy fluxes due to wave-wave interactions. Part I. Methodology and basic results. *J. Fluid Mech.*, **223**, 603–629..
- Shrira, V. I., S. I. Badulin, and C. Khariff, 1996: A model of water wave “horseshoe” patterns. *J. Fluid Mech.*, **318**, 375–404..
- Stiassnie, M., and L. Shemer, 1984: On modification of Zakharov’s equation of surface gravity waves. *J. Fluid Mech.*, **143**, 47–67..
- ,and — ,1987: Energy computations for evolution of class I and II instabilities of Stokes waves. *J. Fluid Mech.*, **174**, 299–312..
- Su, M. Y., 1982a: Three-dimensional deep water waves. Part I. Experimental measurement of skew and symmetric wave patterns. *J. Fluid Mech.*, **124**, 73–108..
- ,1982b: Evolution of groups of gravity waves with moderate to high steepness. *Phys. Fluids*, **25**, 2167–2174..
- ,and W. Green, 1984: Coupling two- and three-dimensional instabilities of surface gravity waves. *Phys. Fluids*, **27**, 2595–2597..
- Young, I. R., L. A. Verhagen, and S. K. Khatri, 1996: The growth of fetch limited waves in water of finite depth. Part 3. Directional spectra. *Coast. Eng.*, **29**, 101–121..
- Zakharov, V. E., 1991: Inverse and direct cascade in the wind-driven surface wave turbulence and wave breaking. *Breaking Waves IUTAM Symposium*, M. L. Banner and R. H. J. Grimshaw, Eds., Sydney, Australia, IUTAM, 69–91..
- ,1998: Weak nonlinear wave-wave interactions in shallow water. *Amer. Math. Soc. Transl.*, **182**, 167–197..

---

## Figures





[Click on thumbnail for full-sized image.](#)

Fig. 1. (a) The nonlinear transfer for five-wave interactions, using JONSWAP input spectrum (with peakedness  $\gamma = 3.3$ ) with Hasselmann–Mitsuyasu spreading as given in [Eqs. \(1\)–\(2\)](#), steepness 0.3 and water depth, 10 m. (b) As in [Fig. 1a](#), assuming Pierson–Moskowitz input spectrum ( $\gamma = 1$ ) with wide  $\cos^2\theta$  spreading.

*Corresponding author address:* Dr. Ray-Qing Lin, Dept. of the Navy, Naval Surface Warfare Center, David Taylor Model Basin, Carderock Division Headquarters, 9500 MacArthur Blvd., West Bethesda, MD 20817-5700.

E-mail: [rlin@wave2.dt.navy.mil](mailto:rlin@wave2.dt.navy.mil)

[top](#) ▲



© 2008 American Meteorological Society [Privacy Policy and Disclaimer](#)  
Headquarters: 45 Beacon Street Boston, MA 02108-3693  
DC Office: 1120 G Street, NW, Suite 800 Washington DC, 20005-3826  
[amsinfo@ametsoc.org](mailto:amsinfo@ametsoc.org) Phone: 617-227-2425 Fax: 617-742-8718  
[Allen Press, Inc.](#) assists in the online publication of AMS journals.