



Abstract View

[Volume 27, Issue 6 \(June 1997\)](#)

Journal of Physical Oceanography

Article: pp. 926–940 | [Full Text](#) | [PDF \(317K\)](#)

The Influence of Stratification on the Inertial Recirculation

Zhengyu Liu

Department of Atmospheric and Oceanic Sciences, University of Wisconsin—Madison, Madison, Wisconsin

(Manuscript received April 29, 1994, in final form November 13, 1996)

DOI: 10.1175/1520-0485(1997)027<0926:TIOSOT>2.0.CO;2

ABSTRACT

A two-layer quasigeostrophic model is used to investigate the influence of stratification on the inertial recirculation in a full basin model. It is found that the barotropic transport of the inertial recirculation is intensified significantly through barotropic–baroclinic interactions in the presence of a shallow thermocline or a strong stratification. Weakly nonlinear theories and numerical experiments show that a strong baroclinic–barotropic interaction intensifies the advection of potential vorticity anomaly toward the inertial recirculation and therefore forces a stronger recirculation. Furthermore, from the potential vorticity point of view, our model recirculations belong to the generalized “modonlike” recirculation (with $dQ/d\psi < 0$). The increased zonal penetration of recirculation cells with stratification is not caused by the internal dynamics of the recirculation cells. Instead, it is caused by the increased advection of potential vorticity anomaly—an external forcing to the recirculation cells.

Options:

- [Create Reference](#)
- [Email this Article](#)
- [Add to MyArchive](#)
- [Search AMS Glossary](#)

Search CrossRef for:

- [Articles Citing This Article](#)

Search Google Scholar for:

- [Zhengyu Liu](#)

