



Abstract View

[Volume 14, Issue 12 \(December 1984\)](#)

Journal of Physical Oceanography

Article: pp. 1875–1892 | [Abstract](#) | [PDF \(1.30M\)](#)

A Model of the Equatorial Sea Surface Temperature Field and Associated Circulation Dynamics

Lewis M. Rothstein

School of Oceanography and Joint Institute for the Study of the Atmosphere and Ocean, University of Washington, Seattle, WA 98195

(Manuscript received January 10, 1984, in final form October 3, 1984)

DOI: 10.1175/1520-0485(1984)014<1875:AMOTES>2.0.CO;2

ABSTRACT

A linear, continuously stratified analytical model of the equatorial ocean is developed and used to simulate the major features of the equatorial sea surface temperature (SST) field and current system. The model is bounded by meridional barriers to flow in the east and west and is forced by an idealized southeast trade wind field. The steady state response is studied here, although the model is time periodic and capable of simulating the annual cycle or a more general time dependence. The surface heat flux forcing is set to zero so that the contribution of the equatorial upwelling field in maintaining the cool equatorial surface water can be isolated.

Many of the observed features of the equatorial SST field and current system are simulated. The model generates cool SST anomalies on the equator in the eastern half of the basin. The cool water core is located to the south of the equator in the easternmost regions of the basin, with the coldest water along the eastern boundary. The latitudinal and longitudinal scales of the model SST anomaly agree with observations. The latitudinal scales are established by a linear combination of the equatorial deformation radii of all model baroclinic modes and not just the first few modes. A southerly wind that crosses the equator along the eastern boundary is necessary for proper simulation of the equatorial SST field.

The model current structure is familiar. The Equatorial Undercurrent is geostrophic and symmetric about the equator in the center of the basin. It is situated between a surface westward flow and deep westward flow. The eastward volume transport associated with the Undercurrent moves southward in the eastern part of the basin and at some depths, the Equatorial and Coastal Undercurrents are continuous. Boundary effects in the eastern region force the Equatorial Undercurrent to bifurcate at a distance of approximately 100 km from the coast. The southerly core is continuous with the model generated Coastal Undercurrent. All current magnitudes agree reasonably well with observational estimates.

The solution techniques are the standard separation of variables but in order to incorporate a heat flux surface boundary condition, special methods are employed. We have termed these “inverse methods” as we need to

Options:

- [Create Reference](#)
- [Email this Article](#)
- [Add to MyArchive](#)
- [Search AMS Glossary](#)

Search CrossRef for:

- [Articles Citing This Article](#)

Search Google Scholar for:

- [Lewis M. Rothstein](#)

eventually solve for a boundary condition (the SST anomaly field).

top ▲



© 2008 American Meteorological Society [Privacy Policy and Disclaimer](#)

Headquarters: 45 Beacon Street Boston, MA 02108-3693

DC Office: 1120 G Street, NW, Suite 800 Washington DC, 20005-3826

amsinfo@ametsoc.org Phone: 617-227-2425 Fax: 617-742-8718

[Allen Press, Inc.](#) assists in the online publication of *AMS* journals.