

Attention: JavaScript deactivated

Please activate JavaScript to be able to use all website functions!



Quick-Li

MyTUM

Roomfinder

Media services

Faculties

Clinics

Awards and honors

Working at TUM

TUM Fan?

Contact & Directions

Research

Faculties

Researchcenters

Excellence Initiative

Current projects

Research Partners

Research promotion

TUM Graduate School

Postdocs

Open Professorships

Studies

Learning

Degree programs

International Students

Advising

Application and Acceptance

Fees and Financial Aid

During your Studies

Completing your Studies

Continuing Education

Contacts

University Life

Student Life

Accommodations

Sports

Music and Arts

Events

Mentoring


Alumni Services

TUM for schools

ABOUT TUM[Our University](#)**News**[Archive](#)[Publications](#)[Faculties](#)[Clinics](#)[Awards and honors](#)[Working at TUM](#)[TUM Fan?](#)[Contact & Directions](#)

Statistical methods improve biological single-cell analysis

Tracing unique cells with mathematics



Fluorescence- in-situ-hybridization shows mRNA-activity in a tissue sample. Blue: low, red: high activity – Image: S. S. Bajikar / University of Virginia, Charlottesville (USA)

23.01.2014, *Research news*

Stem cells can turn into heart cells, skin cells can mutate to cancer cells; even cells of the type exhibit small heterogeneities. Scientists use single-cell analysis to investigate these. But the method is still laborious and considerable inaccuracies conceal smaller effects. Scientists at Technische Universitaet Muenchen (TUM), the Helmholtz Zentrum Muenchen and the University of Virginia (USA) have now found a way to simplify and improve the analysis by mathematical methods.

Each cell in our body is unique. Even cells of the same tissue type that look identical under the microscope differ from each other. To understand how a heart cell can develop from a stem cell, why one beta-cell produces insulin and another not, or why a normal tissue cell suddenly mutates to a cancer cell, scientists have been targeting the activities of individual RNA molecules.

Proteins are constantly being assembled and disassembled in the cell. RNA molecules read blueprints for proteins and initiate their production. In the last few years scientists around the world have developed sequencing methods that are capable of detecting all active RNA molecules in a single cell at a certain time.

At the end of December 2013 the journal Nature Methods declared single-cell sequencing the "Method of the Year." However, analysis of individual cells is extremely challenging. The handling of the cells generates errors and inaccuracies. Smaller differences in gene regulation can be overwhelmed by the statistical "noise."

Easier and more accurate, thanks to statistics

Scientists led by Professor Fabian Theis, Chair of Mathematical modeling of biological systems at the Technische Universitaet Muenchen and director of the Institute of Computational Biology at the Helmholtz Zentrum Muenchen, have now found a way to considerably improve single-cell analysis by applying methods of mathematical modeling.

Instead of just one cell, they took 16-80 samples with ten cells each. "A sample of ten cells is much easier to handle," says Professor Theis. "With ten times the material, the influences of ambient conditions can be markedly suppressed." However, cells with different properties are then distributed randomly on the samples. His collaborator Christiane Fuchs developed statistical methods to still identify the single-cell properties in the mixture of signals.

Combining model and experiment

On the basis of known biological data, Theis and Fuchs modeled the distribution for the case of genes that exhibit two well-defined regulatory states. Together with Kevin A. Janes and Sameer Bajikar at the University of Virginia in Charlottesville (USA), they were able to prove experimentally that with the help of statistical methods single-cell analysis delivers results of higher accuracy than can be achieved through analysis of the same number of single cell samples.

In many cases, several gene actions are triggered by the same factor. Even in such cases, the statistical method can be applied successfully. Fluorescent markers mark different gene activities. The result is a mosaic, which again can be checked to spot whether different cells respond differently to the factor.

The method is so sensitive that it even shows one deviation in 40 otherwise identical cells. The fact that this difference actually is an effect and not a random outlier was proven experimentally.

This work has been funded by the American Cancer Society, the National Institutes of Health, the German Research Foundation (DFG), the German Academic Exchange Service (DAAD), the Pew Scholars Program in the Biomedical Sciences, the David and Lucile Packard Foundation, the National Science Foundation and the European Research Council.

Publication:

Parameterizing cell-to-cell regulatory heterogeneities via stochastic transcriptional profiles
Sameer S. Bajikar, Christiane Fuchs, Andreas Roller, Fabian J. Theis, and Kevin A. Janes
PNAS, Early Edition, 21 Januar 2014, [Doi: 10.1073/pnas.1311647111](https://doi.org/10.1073/pnas.1311647111)

Contact

Prof. Dr. Fabian J. Theis
Technische Universitaet Muenchen
Mathematical modeling of biological systems
Boltzmannstr. 3, 85748 Garching, Germany
[Tel.: +49 89 289 18386](tel:+498928918386) – [E-Mail](mailto:fabian.theis@tum.de)

Helmholtz-Zentrum München
Institute of Computational Biology
Ingolstädter Landstr. 1, 85764 Neuherberg, Germany
[Tel.: +49 89 3187 2211](tel:+498931872211)
[E-Mail](mailto:icb@helmholtz-muenchen.de) - [Internet](http://www.icb.helmholtz-muenchen.de)

[Desk: Dr. Andreas Battenberg](#)

Recommend page

[Back to News](#)

[Jobs](#)

[Site Map](#)

[Feedback](#)

[Media](#)

[Legal Details](#)

[german](#) [english](#)

Find us on: