



## 栗与美国板栗化感作用的比较

李晓娟<sup>1,2</sup>, 王强<sup>2\*</sup>, 倪穗<sup>1\*</sup>, 阮晓<sup>2</sup>, 王永红<sup>2</sup>, 张焕<sup>2</sup>, 王高峰<sup>3</sup>

<sup>1</sup>宁波大学海洋学院, 浙江宁波 315211;

<sup>2</sup>浙江大学宁波理工学院, 浙江宁波 315100;

<sup>3</sup>School of Agricultural, Forest and Environmental Science, Clemson University, Clemson 29631, USA

### Allelopathy comparison between *Castanea mollissima* and *C. dentata*

LI Xiao-Juan<sup>1,2</sup>, WANG Qiang<sup>2\*</sup>, NI Sui<sup>1\*</sup>, RUAN Xiao<sup>2</sup>, WANG Yong-Hong<sup>2</sup>, ZHANG Huan<sup>2</sup>, and Geoff WANG<sup>3</sup>

<sup>1</sup>School of Marine Sciences, Ningbo University, Ningbo, Zhejiang 315211, China;

<sup>2</sup>Ningbo Institute of Technology, Zhejiang University, Ningbo, Zhejiang 315100, China;

<sup>3</sup>School of Agricultural, Forest and Environmental Sciences, Clemson University, Clemson 29631, USA

[摘要](#)

[图/表](#)

[参考文献](#)

[相关文章](#)

[点击分布统计](#)

[下载分布统计](#)

?