

Turkish Journal of Botany

Turkish Journal
of
Botany

The Family Boletaceae s.l. (Excluding Boletus) in the Republic of Macedonia

Mitko KARADELEV¹, Katerina RUSEVSKA¹, Sofche SPASIKOVA²

¹Institute of Biology, Faculty of Natural Science and Mathematics, P.O. Box 162, Arhimedova 5, 1000 Skopje, The Republic of Macedonia

²Macedonian Mycological Society, Arhimedova 5, P.O Box 162, 1000 Skopje,
The Republic of Macedonia

 [Keywords](#)
 [Authors](#)



bot@tubitak.gov.tr

[Scientific Journals Home](#)

[Page](#)

Abstract: This is the first paper on systematic research into Boletaceae s.l. (excluding *Boletus L.*) in the Republic of Macedonia. A total of 31 species belonging to 8 genera are registered in the Republic of Macedonia. Two genera (*Gyrodon* Opat. and *Phylloporus* Quél.) and 8 species (*Chalciporus amarellus* (Quél.) Bataille, *Gyrodon lividus* (Bull.: Fr.) Opat., *Leccinum variicolor* Watling, *Leccinum vulpinum* Watling, *Phylloporus pelletieri* (Lév.) Quél., *Suillus mediterraneensis* (Jacquet. & J.B. Blum) Redeuilh, *Xerocomus ferrugineus* (Schaeff.) Bon, and *Xerocomus rubellus* (Krombh.) Quél.) are reported as new for Macedonia.

Key Words: Fungi, Boletaceae s.l. (excl. *Boletus L.*), reported species, ecology and distribution, Macedonia

Turk. J. Bot., **31**, (2007), 539-550.

Full text: [pdf](#)

Other articles published in the same issue: [Turk. J. Bot., vol.31,iss.6](#).