Turkish Journal of Botany

Turkish Journal

of

Tuncay DIRMENCI
Balikesir University, Faculty of Education, Department of Biology Education, Balikesir - TURKEY
Bayram YILDIZ
Balikesir University, Faculty of Arts & Sciences, Biology Department, Balikesir - TURKEY
Gülendam TÜMEN
Balikesir University, Faculty of Arts & Sciences, Biology Department, Balikesir - TURKEY
Skewwords

Keywords Authors

Abstract: Four Nepeta L. species endemic to East Anatolia, Nepeta baytopii Hedge & Lamond, N. crinita Montbret & Aucher ex Bentham, N. obtusicrena Boiss. & Kotschy ex Hedge and N. sorgerae Hedge & Lamond, are examined. The areas of these species are quite limited. According to "The Red Book of Turkey" the threatened categories of these species are N. baytopii, EN; N. crinita, EN; N. obtusicrena, LR (cd) and N. sorgerae, Lr(cd).

(cu) and N. sorgerae, Li(cu

bot@tubitak.gov.tr Turk. J. Bot., 28, (2004), 221-226.

Full text: pdf

Scientific Journals Home Page Other articles published in the same issue: Turk. J. Bot., vol.28, iss. 1-2.

Key Words: Nepeta, threat categories, endemism