

# Turkish Journal of Zoology

Turkish Journal


of  
Zoology

On New Populations of *Rana holtzi* and *Rana macrocnemis* (Ranidae: Anura)

İbrahim BARAN<sup>1</sup>, Çetin ILGAZ<sup>2</sup>, Yusuf KUMLUTAŞ<sup>1</sup>, Kurtuluş OLGUN<sup>3</sup>, Aziz AVCI<sup>3</sup>, Fatma İRET<sup>1</sup>  
<sup>1</sup>Dokuz Eylül University, Faculty of Education, Department of Biology, 35150 Buca, İzmir - TURKEY

<sup>2</sup>Dokuz Eylül University, Fauna and Flora Research and Application Center, 35150 Buca, İzmir - TURKEY

<sup>3</sup>Adnan Menderes University, Science and Arts Faculty, Department of Biology, 09010 Aydın - TURKEY

 [Keywords](#)  
 [Authors](#)



[zool@tubitak.gov.tr](mailto:zool@tubitak.gov.tr)

[Scientific Journals Home Page](#)

**Abstract:** *Rana holtzi*, which is only known from Karagöl at altitude 2500 m and Çinigöl at altitude 2600 m, was determined from Eğrigöl for the first time at altitude 3000 m. Thus, the distribution range of *R. holtzi* in the Bolkar Mountains was extended. In addition, specimens of *Rana macrocnemis* were recorded from Seviçova (2500 m a.s.l.), located in the Bolkar Mountains near Ereğli (Konya), for the first time. Furthermore, the village of Örtülü (Maden, Elazığ), where specimens of *Rana macrocnemis* were captured, is determined to be southernmost distribution point of *R. macrocnemis*. Mountain frog specimens captured from new localities were compared in terms of significant taxonomical features with regard to the literature. In addition, ecological observations of localities where specimens were captured were determined.

**Key Words:** *Rana holtzi*, *Rana macrocnemis*, new locality

---

Turk. J. Zool., **31**, (2007), 241-247.

Full text: [pdf](#)

Other articles published in the same issue: [Turk. J. Zool., vol.31, iss.3.](#)