

Turkish Journal of Zoology

Turkish Journal

Decapod (Crustacea) Fauna of Saros Bay (Northeastern Aegean Sea)

of
Zoology

A. Suat ATEŞ

Çanakkale Onsekiz Mart University, Fisheries Faculty, 17100 Çanakkale - TURKEY

Tuncer KATAĞAN, Ahmet KOCATAŞ

Ege University, Fisheries Faculty 35100 Bornova, İzmir - TURKEY

Füsun ERKAN YURDABAK

Çanakkale Onsekiz Mart University, Fisheries Faculty, 17100 Çanakkale - TURKEY

 [Keywords](#)
[Authors](#)

Abstract: The specific composition of decapod crustaceans collected in the sublittoral depths (8-135 m) of Saros Bay (northeastern Aegean Sea) is presented. A total of 40 species (17 Natantia, 1 Macrura Reptantia, 8 Anomura and 14 Brachyura) were recorded. The dominant taxon is Natantia, represented by a total of 17 species and an occurrence frequency of 42%.

Key Words: Decapoda Crustacea, Saros Bay, Northeastern Aegean Sea, Turkey



zool@tubitak.gov.tr

Turk. J. Zool., **29**, (2005), 119-124.

Full text: [pdf](#)

[Scientific Journals Home](#)
[Page](#)

Other articles published in the same issue: [Turk. J. Zool., vol.29, iss.2.](#)