

Turkish Journal of Zoology

Turkish Journal

of

Zoology

Feeding Organisms Living in *Capoeta trutta* (Heckel;1843) in the Tigris River and Changes in these Organisms According to Month and Age

Fevzi YILMAZ

Dumlupınar Üniversitesi, Fen Edebiyat Fakültesi, Biyoloji Bölümü, Kütahya - TÜRKİYE

Kemal SOLAK

Gazi Üniversitesi, Gazi Eğitim Fakültesi, Biyoloji Eğitimi Bölümü, Ankara - TÜRKİYE

 [Keywords](#)
[Authors](#)



zool@tubitak.gov.tr

[Scientific Journals Home](#)
[Page](#)

Abstract: This study was conducted in order to determine the species of feeding organisms living in the stomach contents of *Capoeta trutta* in the Tigris River and changes in the organisms according to season, month and age. For this purpose, 120 fish samples were caught between September 1991 and August 1992. At the end of this study, 31 plant-like and 2 animal-like organisms were identified, making a total of 33 genera. The total nutrition consisted of the following percentages: 61.24% Bacillariophyta, 27.72% Cyanophyta, 8.88% Chlorophyta, 1.88% Euglenophyta and 0.25% zoological organisms.

Key Words: *Capoeta trutta*, food types, Tigris River

Turk. J. Zool., **23**, (1999), 973-978.

Full text: [pdf](#)

Other articles published in the same issue: [Turk. J. Zool., vol.23,iss.EK3](#).