



Physics > Fluid Dynamics

# Efficiency of size-dependent particle separation by pinched flow fractionation

Aparna Srivastav, Thomas Podgorski, Gwennou Coupier

(Submitted on 14 Apr 2012)

Pinched flow fractionation is shown to be an efficient and selective way to quickly separate particles by size in a very polydisperse semi-concentrated suspension. In an effort to optimize the method, we discuss the quantitative influence of the pinching intensity in the balance between the requirements of selectivity and minimal dilution.

Comments: 5 pages, 3 figures, accepted in Microfluidics and Nanofluidics

Subjects: **Fluid Dynamics (physics.flu-dyn)**; Soft Condensed Matter (cond-mat.soft)

Cite as: **arXiv:1204.3153v1** [physics.flu-dyn]

## Submission history

From: Gwennou Coupier [[view email](#)]

[v1] Sat, 14 Apr 2012 08:20:03 GMT (142kb,D)

*[Which authors of this paper are endorsers?](#)*

Link back to: [arXiv](#), [form interface](#), [contact](#).

## Download:

- [PDF](#)
- [Other formats](#)

Current browse context:

[physics.flu-dyn](#)

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1204](#)

Change to browse by:

[cond-mat](#)

[cond-mat.soft](#)

[physics](#)

## References & Citations

- [NASA ADS](#)

## Bookmark (what is this?)

