



Nuclear Theory

Spectra, flow and HBT in Pb-Pb collisions at the LHC

[Piotr Bozek](#)

(Submitted on 29 Jun 2011)

The transverse momentum spectra, elliptic flow and interferometry radii for Pb-Pb collisions at the LHC are calculated in relativistic viscous hydrodynamics. For Glauber model initial conditions, we find that the data can be described using a small value of shear viscosity $\eta/s=0.08$. The viscosities and the equation of state are the same as used for RHIC energies.

Comments: To appear in the conference proceedings for Quark Matter 2011, May 23 - May 28, Annecy, France

Subjects: **Nuclear Theory (nucl-th)**; Nuclear Experiment (nucl-ex)

Journal reference: J. Phys. G: Nucl. Part. Phys. 38 (2011) 124043

DOI: [10.1088/0954-3899/38/12/124043](https://doi.org/10.1088/0954-3899/38/12/124043)

Cite as: [arXiv:1106.5953](https://arxiv.org/abs/1106.5953) [nucl-th]
(or [arXiv:1106.5953v1](https://arxiv.org/abs/1106.5953v1) [nucl-th] for this version)

Submission history

From: Bozek [[view email](#)]

[v1] Wed, 29 Jun 2011 14:43:18 GMT (17kb)

[Which authors of this paper are endorsers?](#)

Link back to: [arXiv](#), [form interface](#), [contact](#).

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

nucl-th

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1106](#)

Change to browse by:

[nucl-ex](#)

References & Citations

- [INSPIRE HEP](#)
([refers to](#) | [cited by](#))
- [NASA ADS](#)

Bookmark([what is this?](#))

