化学

核径迹二甲基甲酰胺溶液敏化特性、机理及新现象研究

王玉兰,徐世平

清华大学 核能与新能源技术研究院, 北京 102201

收稿日期 2008-1-23 修回日期 2008-3-25 网络版发布日期: 2008-7-1

摘要 实验研究了二甲基甲酰胺溶液敏化剂对PET材料中核径迹蚀刻特性改变的规律,描述了实验中观察到的新现象,并初步分析了敏化作用机理。

关键词 二甲基甲酰胺溶液; 敏化; 核径迹

分类号 TL815

New Phenomenon and Dimethyl Formamide Treatment Ef fect

in Polymer Irradiated With Heavy Ions

WANG Yu-lan, XU Shi-ping

Institute of Nuclear and New Energy Technology, Tsinghua University, P.O.Box 1021, Beijing 102201, China

Abstract The dimethyl formamide (DMF) treatment effect in PET irradiated with 142 MeV ³ ²S was investigated. It is found that the change of track etching rate with changing the time and te mperature of DMF treatment, with changing the time dipping in water before etching and changin g the time stored in air after DMF treatment. New phenomenon observed in polymer dipped in w ater for more than 10 min between DMF treatment and etching process is also described in the p aper.

Key words dimethyl formamide solvent sensibilization nuclear track

扩展功能

本文信息

- ► Supporting info
- ▶ [PDF全文](3348KB)
- ▶[HTML全文](0KB)
- ▶参考文献

服务与反馈

- ▶把本文推荐给朋友
- ▶文章反馈
- ▶浏览反馈信息

相关信息

- ▶ <u>本刊中 包含"二甲基甲酰胺溶液;</u> 敏化;核径迹"的 相关文章
- ▶本文作者相关文章
- 王玉兰
- ・ 徐世平

通讯作者