



# Agricultural Journals

*Czech Journal of*

**FOOD SCIENCE**

[home](#) [page](#) [about us](#) [contact](#)

[us](#)

## Table of Contents

**IN PRESS**

**CJFS 2014**

**CJFS 2013**

**CJFS 2012**

**CJFS 2011**

**CJFS 2010**

**CJFS 2009**

**CJFS 2008**

**CJFS 2007**

**CJFS 2006**

**CJFS 2005**

**CJFS 2004**

**CJFS 2003**

**CJFS 2002**

**CJFS 2001**

**CJFS Home**

## **Editorial Board**

### **For Authors**

- **Authors Declaration**
- **Instruction to Authors**
- **Guide for Authors**
- **Copyright Statement**
- **Submission**

### **For Reviewers**

- **Guide for Reviewers**
- **Reviewers Login**

---

### **Subscription**

# **Czech J. Food Sci.**

**Atanassova J.,  
YurukovaL., Lazarova**

# Pollen and inorganic characteristics of Bulgarian unifloral honeys

Czech J. Food Sci., 30 (2012): 520-526

The melissopalynological characteristics three main physicochemical parameters (water content, pH, and electrical conductivity), and 19 macro- and microelements contents of 15 honey types from throughout Bulgaria that were collected from 2006 to 2009 were evaluated. The main honeys studied came from *Robinia pseudoacacia* L., *Helianthus annuus* L., *Brassica* spp., *Tilia* spp., and *Vicia* spp. The botanical origins of unifloral honey samples were identified as *Lotus* spp., *Coriandrum sativum* L., *Daucus*-type, *Stachys*-type *Salix* spp., *Prunus* spp., *Castanea sativa* Mill., *Paliurus spina-christi* Mill., *Sophora japonica*, and *Amorpha* spp. Based on the physicochemical parameters and