

Exopolysaccharide Production by *Streptococcus salivarius* ssp. *thermophilus* Cultures. 1. Conditions of Production

Frederique Gancel¹ and Georges Novel¹

¹ Laboratoire de Génétique Microbienne, Institut de Recherche en Biologie Appliquée, Université de Caen, 14032 Caen, Cedex, France

An industrial strain of *Streptococcus salivarius* ssp. *thermophilus*, grown in a semisynthetic medium, produces EPS during the stationary phase of growth. Temperatures or sugars that decrease growth rate increase polymer synthesis. Factorial experiments showed that the nature of the sugar, temperature, and initial pH interfere with polymer production. At optimal growth rate, polysaccharide production is dependent on lactose.

Key Words: exopolysaccharide • *Streptococcus salivarius* ssp. *thermophilus* • lactose • sucrose

Submitted on March 4, 1993

Accepted on October 18, 1993

This Article

- ▶ [Full Text \(PDF\)](#)
- ▶ [Alert me when this article is cited](#)
- ▶ [Alert me if a correction is posted](#)

Services

- ▶ [Similar articles in this journal](#)
- ▶ [Alert me to new issues of the journal](#)
- ▶ [Download to citation manager](#)
- ▶ [Get Permissions](#)

Citing Articles

- ▶ [Citing Articles via HighWire](#)
- ▶ [Citing Articles via Google Scholar](#)

Google Scholar

- ▶ [Articles by Gancel, F.](#)
- ▶ [Articles by Novel, G.](#)
- ▶ [Search for Related Content](#)

PubMed

- ▶ [Articles by Gancel, F.](#)
- ▶ [Articles by Novel, G.](#)

This article has been cited by other articles:



Journal of Dairy Science

▶ HOME

G. Robitaille, S. Moineau, D. St-Gelais, C. Vadeboncoeur, and M. Britten
Detection and quantification of capsular exopolysaccharides from *Streptococcus thermophilus* using lectin probes.

J Dairy Sci, November 1, 2006; 89(11): 4156 - 4162.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)



Journal of Dairy Science

▶ HOME

P. Ruas-Madiedo and C. G. de los Reyes-Gavilan
Invited Review: Methods for the Screening, Isolation, and Characterization of Exopolysaccharides Produced by Lactic Acid Bacteria

J Dairy Sci, March 1, 2005; 88(3): 843 - 856.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)




E. P. Briczinski and R. F. Roberts
Production of an Exopolysaccharide-Containing Whey Protein Concentrate by Fermentation of Whey
J Dairy Sci, December 1, 2002; 85(12): 3189 - 3197.
[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)




Applied and Environmental Microbiology

P. L. Pham, I. Dupont, D. Roy, G. Lapointe, and J. Cerning
Production of Exopolysaccharide by *Lactobacillus rhamnosus* R and Analysis of Its Enzymatic Degradation during Prolonged Fermentation
Appl. Envir. Microbiol., June 1, 2000; 66(6): 2302 - 2310.
[\[Abstract\]](#) [\[Full Text\]](#)



Applied and Environmental Microbiology

P. J. Looijesteijn, I. C. Boels, M. Kleerebezem, and J. Hugenholtz
Regulation of Exopolysaccharide Production by *Lactococcus lactis* subsp. *cremoris* by the Sugar Source
Appl. Envir. Microbiol., November 1, 1999; 65(11): 5003 - 5008.
[\[Abstract\]](#) [\[Full Text\]](#)



Applied and Environmental Microbiology

B. Degeest and L. De Vuyst
Indication that the Nitrogen Source Influences Both Amount and Size of Exopolysaccharides Produced by *Streptococcus thermophilus* LY03 and Modelling of the Bacterial Growth and Exopolysaccharide Production in a Complex Medium
Appl. Envir. Microbiol., July 1, 1999; 65(7): 2863 - 2870.
[\[Abstract\]](#) [\[Full Text\]](#)



Applied and Environmental Microbiology

S. A. Kimmel, R. F. Roberts, and G. R. Ziegler
Optimization of Exopolysaccharide Production by *Lactobacillus delbrueckii* subsp. *bulgaricus* RR Grown in a Semidefined Medium
Appl. Envir. Microbiol., February 1, 1998; 64(2): 659 - 664.
[\[Abstract\]](#) [\[Full Text\]](#)