Scientific Research Open Access



Search Keywords, Title, Author, ISBN, ISSN

	Home Journals Books Conferences News	About Us	s Job	
ŝ	Home > Journal > Earth & Environmental Sciences > JEP	Open Spec	ial Issues	
Inde	xing View Papers Aims & Scope Editorial Board Guideline Article Processing Charges	Published Special Issues		
JEP>	IEP> Vol.1 No.1, March 2010		Special Issues Guideline	
Polyprenol from the Whole Plants of Leucaena leucocephala		JEP Subscription		
	PDF (Size:80KB) PP. 70-72 DOI: 10.4236/jep.2010.11009 Author(s) Chung-Yi Chen, Yau-Der Wang		Most popular papers in JEP	
			About JEP News	
	ABSTRACT icaprenol-11 (polyprenol) (1), squalene (2), lupeol (3), ?-sitostenone (4), trans-coumaric acid (5), cis- oumaric acid (6) pheophytin-a (7), pheophorbide a methyl ester (8), methyl-132-hydroxy-(132-S)- heophorbide-b (9) and aristo-phyll-C (10) were isolated from the whole plants of Leucaena leucocephala Leguminosae). Among them, 1 and 2 was found for the first time from this species. The structure of these ompounds were characterized and identified by spectra analyses.		Frequently Asked Questions	
cour			Recommend to Peers	
. 0			Recommend to Library	
	WORDS prenol, Leucaena leucocephala, Leguminosae	Contact Us		
Cite this paper C. Chen and Y. Wang, "Polyprenol from the Whole Plants of Leucaena leucocephala," <i>Journal of</i> <i>Environmental Protection</i> , Vol. 1 No. 1, 2010, pp. 70-72. doi: 10.4236/jep.2010.11009.		Downloads:	301,518	
		Visits:	674,187	
Refe	Perences A. M. Gamal-Eldeen, H. Amer, W. A. Helmy, R. M. Talaat, and H. Ragab, " Chemically-modified polysaccharide extract derived from Leucaena leucocephala alters Raw 264.7 murine macrophage functions," International Immuno-phar- macology, Vol. 7, pp. 871-878, February 2007.	Sponsors, Associates, a Links >>		
[2]	A. J. Erickson, R. S. Ramsewak, A. J. Smucker, and M. G. Nair, "Nitrification Inhibitors from the roots of Leucaena leucocephala," Journal of Agricultural and Food Chemistry, Vol. 48, pp. 6174-6177, December 2000.	 The International Conference of Pollution and Treatment Technology (PTT 2013) 		
[3]	T. D. Xuan, A. A. Elzaawely, F. Deba, M. Fukuta, and S. Tawata, "Mimosine in Leucaena leucocephala as a potent bio-herbicide," Agronomy for Sustainable Development, Vol. 26, pp. 89-97, April 2006.			
[4]	P. H. Kaufusi, L. S. Forsberg, P. Tittabutr, and D. Borthakur, "Regulation of exopolysaccharide synthesis in rhizobium sp. strain TAL1145 involves an alternative sigma factor gene, rpoH2," Microbiology, Vol. 150, pp. 3473-3482, October 2004.			
[5]	P. K. K. Yeung, F. T. W. Wong, and J. T. Y. Wong, "Mimosine, the allelochemical from the leguminous tree Leucaena leucocephala, selectively enhances cell prolix- feration in dinoflagellates," Applied and Environmental Micro-biology, Vol. 68, pp. 5160-5163, October 2002.			
[6]	M. Soedarjo, and D. Borthakur, "Mimosine, a toxin produ- ced by the tree-legume leucaena provides competition advantage to mimosine-degrading rhizobium strains," Soil Biology and Biochemistry, Vol. 30, pp. 1605-1613, October 1998.			
[7]	Y. Kamada, N. Oshiro, M. Miyagi, H. Oku, F. Hongo, and I. Chinen, "Osteopathy in broiler chicks fed toxic mimosine in Leucaena leucocephala," Bioscience Biotechnology and Biochemistry, Vol. 62, pp. 34-38, January 1998.			
[8]	I. M. Villasenor, R. M. T. Gajo, and R. C. Gonda, "Bio- activity studies on the alkaloid extracts from seeds of Leucaena leucocephala," Phytotherapy Research, Vol. 11, pp. 615-617, December 1998.			

[9] I. L. Tsai, Y. F. Jeng, C. Y. Duh, and I. S. Chen, " Cytotoxic constituents from the leaves of Litsea akoensis," The Chinese Pharmaceutical Journal, Vol. 53, pp. 291-301, December 2001.

- [10] K. Seki, T. Tomihari, K. Haga, and R. Kaneko, "Iristec- torenes A and C-G, monocyclic triperpene esters from Iris tectorum," Phytochemistry, Vol. 53, pp. 425-431, May 1994.
- [11] C. Zhao, J. Shao, and X. Li, " Chemical constituents from fruits of Ailanthus altissima," Zhongguo Zhong Yao Za Zhi, Vol. 34, pp. 2197-2199, September 2009.
- [12] C. Y. Jiang, S. Z. Mu, B. Deng, Y. H. Ge, J. X. Zhang, and X. J. Hao, " Studies on the chemical constituents from Euphorbia chrysocoma," Zhong Yao Cai, Vol. 32, pp. 1390-1392, September 2009.
- [13] C. J. Li, A. A. Ahmed, A. C. Arias, and T. J. Mabry, " Clerodane diterpenoids, long chain esters of coumaric acid and other compounds from Baccharis myrsinites," Phytochemistry, Vol. 45, pp. 571-574, June 1997.
- [14] W. T. Li, H. W. Tsao, Y. Y. Chen, S. W. Cheng, and Y. C. Hsu, " A study on the photodynamic properties of chlorophyll derivatives using human hepatocellular carcinoma cells," Photochemical and Photobiological Sciences, Vol. 6, pp. 1341-1348, December 2007.