Scientific Research



Search Keywords, Title, Author, ISBN, ISSN

Technology (PTT 2013)

Home	Journals	Books	Conferences	News	About Us	Jobs
Home > Journal > Earth & Environmental Sciences > JEP					Open Special Issues	
Indexing View Papers Aims & Scope Editorial Board Guideline Article Processing Charges					Published Special Issues	
JEP> Vol.3 No.11, November 2012					Special Issues Guideline	
OPEN©ACCESS Comparison of the Groundwater Quality in the West Tahta Area,					JEP Subscription	
Upper Egypt in 1989 and 2011 PDF (Size: 1821KB) PP. 1442-1457 DOI: 10.4236/jep.2012.311162					Most popular papers in JEP	
Author(s)					About JEP News	
Ismail Esam, Faid Abdalla, Niesner Erich, Mauritsch Hermann ABSTRACT					Frequently Asked Questions	
Egypt is characterized by arid climate with rare rainfall that means surface water. The River Nile is the main source of water supply. The study area is located in Upper Egypt in the western desert. The area under					Recommend to Peers	
investigation is subjected to intensive development in the last twenty years, clearly seen in the change of the agriculture and urban areas. The rapid increase of the population in Egypt especially in the upper part					Recommend to Library	
leads to an increasing demand for houses on the old cultivated land lead to the migration towards the desert and reclaimed new areas. These new reclaimed areas are depending on the ground water. The extensive use of groundwater and increasing fertilizing system leads to the change of the water table,					Contact Us	
characterization a	nd quality. In this stud	y we use the analysis	of 30 groundwater san analyzed in 2011 from	nples collected and	Downloads:	301,489
identify the change	ntify the change in the quality of the ground water. According the hydrogeologic data which were lected from the study area in 1989 and 2011 we found a changing water table decreasing, from 1 to 48				Visits:	672,811
m in 1989 and 5 to 60 m in 2011. The water level is decreasing in the same direction towards the River Nile. The sufficient recharge and leaching and dissolution of the aquifer materials due to extensive use lead to increase in TDS which reaching to 4453 ppm in 2011. The water type in the 1989 samples is mainly NaHCO <sub>3</sub>					Sponsors, Associates, a Links >>	
and in 2011 samples $Na_2SO_4$ . Grid classification shows that the water samples collected in 1989 are mainly sodium bicarbonate and sodium chloride extending to sodium sulphate while the water samples collected in 2011 are mainly sodium sulphate and sodium chloride. Most of the 1989 groundwater samples are suitable					The International Conference of Pollution and Treatment	

**KEYWORDS** 

Under Groundwater; Withdrawal; West Tahta; Evaluation of Groundwater

## Cite this paper

I. Esam, F. Abdalla, N. Erich and M. Hermann, "Comparison of the Groundwater Quality in the West Tahta Area, Upper Egypt in 1989 and 2011," *Journal of Environmental Protection*, Vol. 3 No. 11, 2012, pp. 1442-1457. doi: 10.4236/jep.2012.311162.

for drinking purposes while majority of the 2011 groundwater samples are not suitable due to high salinity.

## References

- RIGW & IWACO, " Groundwater Development for Irrigation and Drainage in the Nile Valley," Groundwater Development in the Area of West Tahta, Technical Note 70, Internal Report 124-89-05, 1989.
- [2] R. Said, "The Geologic Evaluation of the River Nile," Springer-Verlag, New York, 1981, p. 151.
- [3] R. Said, " The Geology of Egypt," Balkema Publisher, Rotterdam, 1990.
- [4] Meteorological Authority of Egypt (MAE), " Meteorological Database, Cairo, Egypt," 2000.
- [5] A. A. Ayman, "Using Generic and Pesticide DRASTIC GIS-Based Models for Vulnerability Assessment of the Quaternary Aquifer at Sohag, Egypt," Hydrogeology Journal, Vol. 17, No. 5, 2009, pp. 1203-1217. doi:10.1007/s10040-009-0433-3
- [6] A. A. Omran, E. A. Korany and A. A. Abdel-Rahman, " An Integrated Approach to Evaluate Groundwater Potentiality—A Case Study," Proceeding of the 3rd International Conference of Applied

Geophysical, 18-20 March 2006, Cairo.

- [7] A. O. Awad, "Integration of Remote Sensing, Geophysics and GIS to Evaluate Groundwater Potentiality—A Case Study in Sohag Region," Egypt, 2008. http://faculty.ksu.edu.sa/72005/Papers%20of%20Interest%20Remote%20Sensing/Integration% 20of%20Remote%20Sensing,%20Geophysics%20and%20Gis.pdf
- [8] J. D. Hem, " Study and Interpretation of the Chemical Characteristics of Natural Water," US Geological Survey, Water Supply Paper 2254, 1989, p. 264. http://pubs.usgs.gov/wsp/wsp2254/
- [9] J. D Hem, " Study and Interpretation of the Chemical Characteristics of Natural Water," US Geological Survey, Water Supply Paper 2254, 1985, pp.117, 120, 264.
- [10] H. M. Ragunath, " Groundwater," Geological Journal, Vol. 18, No. 3, 1987, pp. 279-280.
- [11] D. K. Todd, " Groundwater Hydrology," Wiley, New York, 1980, p. 315.
- [12] G. N. Sawyer and D. L. McCarthy, " Chemistry of Sanitary Engineers," McGraw Hill, New York, 1967, p. 518.
- [13] A. M. Piper, " A Graphic Procedure in the Geochemical Interpretation and Analysis of Water Samples," US Geological Survey, Water Supply Paper 1454, 1944.