

[Available Issues](#) | [Japanese](#)>> [Publisher Site](#)Author: [ADVANCED](#)

Volume Page

Keyword: [TOP](#) > [Available Issues](#) > [Table of Contents](#) > [Abstract](#)

ONLINE ISSN : 1349-1008

PRINT ISSN : 1343-943X

Plant Production Science

Vol. 11 (2008) , No. 3 352-354

[\[PDF \(530K\)\]](#) [\[References\]](#)**Effects of Shading on Hydraulic Resistance and Morphological Traits of Internode and Node of Napiergrass (*Pennisetum purpureum* Schumach.)**[Kiyoshi Nagasuga](#)¹⁾ and [Fumitake Kubota](#)¹⁾

1) Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University

(Received: February 9, 2007)

Keywords: [Cross-sectional area](#), [Hydraulic resistance](#), [Internode](#), [Internode length](#), [Napiergrass](#), [Shade](#)[\[PDF \(530K\)\]](#) [\[References\]](#)Download Meta of Article[\[Help\]](#)[RIS](#)[BibTeX](#)

To cite this article:

Kiyoshi Nagasuga and Fumitake Kubota: "Effects of Shading on Hydraulic Resistance and Morphological Traits of Internode and Node of Napiergrass (*Pennisetum purpureum* Schumach.)". *Plant Production Science*, Vol. **11**, pp.352-354 (2008) .

doi:10.1626/pps.11.352

JOI JST.JSTAGE/pps/11.352

Copyright (c) 2008 by The Crop Science Society of Japan



[Japan Science and Technology Information Aggregator, Electronic](#)

