



[Available Issues](#) | [Japanese](#)

Author:  [ADVANCED](#) | Volume  Page

Keyword:



[TOP](#) > [Available Issues](#) > [Table of Contents](#) > **Abstract**

## Horticultural Research (Japan)

Vol. 8 (2009) , No. 4 427-432

[E]

### Nursery-grafting Method for Persimmon Propagation Rootstock

[Masaru Wakisaka](#)<sup>1)</sup>, [Teruhiko Sugimura](#)<sup>1)</sup>, [Tomoya Ishimori](#)<sup>2)</sup> and

1) Nara Fruit Tree Research Center

2) Laboratory of Horticultural Science. Faculty of Agriculture, Kinki University

(Received September 26, 2008)

(Accepted March 13, 2009)

We present a new nursery grafting method for persimmon (*Diospyros*) propagation. This study examined the effects of grafting time and leaf graft-take ratio and growth of grafted plants. Between April and September to graft a scion of 'Fuyu' persimmon onto a current seedling of 'Hokuriku' which was between 110 and 195 days old. Although grafting onto a seedling without leaves yielded a low graft-take ratio, 80 to 100% of scions with leaves were established successfully. Between rootstock with a

significant difference was noted in the callus-forming ratios 20 days : was observed in all nursery plants grafted onto rootstock with leave bud-break feasibility.

**Key Words:** [callus](#), [graft-take ratio](#), [grafting time](#), [leaves](#)

[\[PDF \(1194K\)\]](#) [\[References\]](#)

Downlo

To cite this article:

Masaru Wakisaka, Teruhiko Sugimura, Tomoya Ishimori and Shir  
Nursery-grafting Method for Persimmon Propagation Using a Cur  
(Japan) 8: 427-432 .

---

doi:10.2503/hrj.8.427