Czech Academy of Agricultural Sciences



Open Access Agricultural Journals

Journal of

FOREST SCIENCE

home page about us contact

US

Table of Contents

IN PRESS

JFS 2015

JFS 2014

JFS 2013

JFS 2012

JFS 2011

JFS 2010

JFS 2009

JFS 2008

JFS 2007

JFS 2006

JFS 2005

JFS 2003 JFS Home

Editorial Board

For Authors

- AuthorsDeclaration
- Instruction to Authors
- Guide for Authors
- CopyrightStatement
- Submission

For Reviewers

- Guide for Reviewers
- ReviewersLogin

Subscription

Journal of Forest Science

Root system development and health condition of sycamore maple (*Acer pseudoplatanus* L.) in the air-polluted region of Krušné hory Mts.

Mauer O., Pop M., Palátová E.:

J. For. Sci., 53 (2007): 452-461

[fulltext]

The paper presents results from a study of sycamore maple development, health condition and growth in forest altitudinal vegetation zones (FAVZ) 6 and 7 occurring in pollution damage zones A and B in the air-polluted region of Krušné hory Mts. as compared with the trees of identical height in FAVZ 4 and 5 occurring in pollution damage zone D in the Bohemian-Moravian Upland. Sycamore maple develops a fully diversified root system. On spread mounds it creates only a superficial root system and its growth is retarded. The growth of sycamore maple is limited by the layer of humus horizons. If the layer thickness is over 20 cm, the sycamore roots would grow into mineral horizons.

Keywords:

sycamore maple; root system; humus; afforestation; mounds

[fulltext]

© 2015 Czech Academy of Agricultural Sciences



