



Table of Contents

In Press

Article Archive

JFS (64) 2018

JFS (63) 2017

JFS (62) 2016

JFS (61) 2015

JFS (60) 2014

JFS (59) 2013

JFS (58) 2012

JFS (57) 2011

JFS (56) 2010

JFS (55) 2009

Issue No. 1 (1-50)

Issue No. 2 (51-100)

Issue No. 3 (101-144)

Issue No. 4 (144-192)

Issue No. 5 (194-250)

Issue No. 6 (251-298)

Issue No. 7 (299-344)

Issue No. 8 (345-394)

Issue No. 9 (395-436)

Issue No. 10 (437-483)

Issue No. 11 (485-531)

Issue No. 12 (533-590)

JFS (54) 2008

JFS (53) 2007

JFS (52) 2006

JFS (51) 2005

JFS (50) 2004

JFS (49) 2003

Editorial Board

Ethical Standards

Peer Review Process

Reviewers 2017

For Authors

Author Declaration

Instruction for Authors

Submission Templates

Guide for Authors

Copyright Statement

Submission/Login

The first reaction of soil mite fauna (Acari, Mesostigmata) caused by conversion of Norway spruce stand in the Szklarska Poręba Forest District

M. Skorupski, G. Butkiewicz, A. Wierzbicka

<https://doi.org/10.17221/19/2009-JFS>

Citation: Skorupski M., Butkiewicz G., Wierzbicka A. (2009): The first reaction of soil mite fauna (Acari, Mesostigmata) caused by conversion of Norway spruce stand in the Szklarska Poręba Forest District. *J. For. Sci.*, 55: 235-243.

[download PDF](#)

To observe the first reaction of Mesostigmata mites for conversion of even aged spruce stands under conversion with common beech and silver fir, four pairs of experimental plots directly neighbouring each other were examined. On the spruce stands that started to be converted into beech or fir stands a few years ago, some significant composition and quantitative changes can be observed as regards Mesostigmata order mites. The first reaction of Mesostigmata mites after starting of forest conversion was an increasing number of mite species in the plots. On the plots with relatively close occurrence of beech, increasing number of specimens was also observed in plots under conversion. The share of Parasitidae family mites was decreased on the converted plots with relatively close occurrence of beech. On the converted spruce stands the percentage share of Uropodina suborder mites was found to increase or remain high.

Keywords:

Acari; Mesostigmata; mites; forest conversion; biodiversity; soil ecology

[download PDF](#)

SJR (SCImago Journal Rank – SCOPUS)

2017: 0.206 – Q4 (Forestry)



Share

New Issue Alert

Join the journal on [Facebook!](#)
Ask for [email notification](#).

Publish with JFS!

- Full Open Access
- Rapid review and fast publication
- International knowledge sharing
- No article processing charge

Similarity Check

All the submitted manuscripts are checked by the [CrossRef Similarity Check](#).

Referred to in

- Agrindex of AGRIS/FAO database
- CAB Abstracts
- CNKI
- Czech Agricultural and Food Bibliography
- DOAJ (Directory of Open Access Journals)
- Elsevier's Bibliographic Databases
- Google Scholar
- J-Gate
- SCOPUS
- TOXLINE PLUS
- Web of Science (BIOSIS Citation Index)

Licence terms

All content is made freely available for non-commercial purposes, users are allowed to copy and redistribute the material, transform, and build upon the material as long as they cite the source.

Open Access Policy

This journal provides immediate open access to its content on the principle that making research freely available to the public supports a greater global exchange of knowledge.

Contact

Mgr. Petra Kolářová
Executive Editor
phone: + 420 227 010 355
e-mail: jfs@caav.cz

Address

Journal of Forest Science
Czech Academy of Agricultural Sciences

For Reviewers

Slezská 7, 120 00 Praha 2, Czech
Republic

[Guide for Reviewers](#)

[Reviewers Login](#)

[Subscription](#)

© 2018 Czech Academy of Agricultural Sciences