Czech Academy of Agricultural Sciences Open Access Agricultural Journal VETERINÁRNÍ MEDICÍ VETME age about us contact us **Table of Contents VETMED** 2015 VETMED 2014 VETMED 2013 **VETMED** 2012 **VETMED** 2011 **VETMED** 2010

VETMED 2009

VETMED 2008 **VETMED** 2007 **VETMED** 2006 **VETMED** 2005 **VETMED** 2004 **VETMED** 2003 **VETMED** 2002 **VETMED** 2001 **VETMED** Home

Editorial Board

For Authors

- Authors
 Declaration
- Instruction to Authors
- Guide for

Authors

- Fees
- Submission

Subscription

Veterinarni Medicina

Evaluation of the functional capacity in dogs with naturally acquired heart disease

Agudelo CF, Schanilec P:

Veterinarni Medicina, 58 (2013): 264-27([fulltext]

Functional capacity is the physical ability to perform a defined task. In humans it can be evaluated by using exercise testing, for instance sub-maximal exercis tests. Walking tests are widely used variations of sub-maximal exercise tests and in human medicine are considered to be realistically related to daily physical activity and prognosis in patients with heart failure. The aim of this study was to assess a sub-maximal exercise test in dogs with varying degrees of heart disease. The 6-minute walking test was used to test the functional capacity in healthy dogs and dogs with heart disease (degenerative mitral valve disease). Three groups of untrained owned dogs

were compared. Two groups were dogs had mild and moderate degenerative mitral valve disease (ME-1 and ME-2, respectively) and the third group comprised healthy dogs with the same age and size conditions. Both groups of dogs with mitral valve disease walked shorter distances during 6 min (control 448.92 m; ME-1: 406.89 m; ME-2: 350.04 m). The influence of the test on the heart rate, electrocardiography and blood pressure was also evaluated. Although the 6-minute walking test has already been tested in healthy dogs, dog with chronic pulmonary disease and dog with artificially induced heart disease, this is the first time that this test has been