

Repeated administration of different hormonal preparations for artificial propagation and their effects on reproduction, survival and blood biochemistry profiles of female tench (*Tinca tinca* L.)

J. KOUŘIL¹, M. SVOBODA², J. HAMÁČKOVÁ¹, P. KALÁB², J. KOLÁŘOVÁ¹, A. LEPIČOVÁ¹, M. SEDOVA³, L. SAVINA⁴, P. MORENO RENDÓN⁵, Z. SVOBODOVÁ^{1,2}, T. BARTH⁶, B. VYKUSOVÁ¹

¹Research Institute of Fish Culture and Hydrobiology, University of South Bohemia in České Budějovice, Vodňany, Czech Republic

²University of Veterinary and Pharmaceutical Sciences, Brno, Czech Republic

³Russian Federal Research Institute of Fisheries and Oceanography, Moscow, Russia

⁴Department of Ichthyology and Hydrobiology, Kaliningrad State University, Kaliningrad, Russia

⁵Physic Department, Ecology Area, Extramadura University, Badajoz, Spain

⁶Institute of Organic Chemistry and Biochemistry Czech Academy of Sciences, Prague, Czech Republic

ABSTRACT: The aim of the study was to compare physiological statuses of stripped and not stripped tench females with ovulation induced by GnRHa or carp pituitary extract (CPE). The comparison was based on selected biochemical stress indices (cortisol, glucose), condition (TP, TGA), and cell membrane characteristics (ALT, AST, CK). In the first and in the second year, 37 and 29 tench females were monitored, respectively. No differences in the values of biochemical parameters in either period were found between the two groups of stripped female tench whose ovulation was induced by GnRHa and CPE, respectively. No significant differences in TP, TGA, ALT, AST and CK values were found between the groups of stripped and not stripped females with ovulation induced by GnRHa or CPE. Different values were found in stress indices, specifically in glucose concentrations. A significant increase ($P < 0.01$) in glucose concentrations was found in tench females immediately after stripping and, in not stripped females, about 48 hrs after ovulation induction. Both artificial stripping and the inability to release eggs are important stress factors for female tench.

Keywords: tench; artificial propagation; hormonal stimulation; haematology; biochemistry profile

The reproduction of the majority of cyprinid fish used in aquaculture usually relies on hormonally induced stripping. In carp, Asian herbivorous species and a number of other fish species, the method based on injecting two separate doses of carp pituitary extract (CPE) has been used for decades. The

use of carp pituitary in artificial propagation of tench was first described by Pokorný (1974). Based on his own experiments, Kouřil (1987) proposed to use a single dose of carp pituitary extract (CPE) in tench and to decrease the originally recommended higher doses to 2 mg/kg. For Asian herbivorous fish

Supported by Czech Research Projects (MSM No. 6007665809) Biological, Environmental and Breeding Aspects of Fishery and (ME 853) Kontakt.

species and salmon, artificial propagation methods based on the use of gonadotropin-releasing hormone (GnRH) for ovulation induction have been developed (Cooperative team, 1977; Donaldson et al., 1981). A possibility of using mammalian GnRH instead of CPE for artificial propagation has also been confirmed in female tench (Kouřil and Barth, 1981; Kouřil et al., 1986). The authors ascertained that the administration of mammalian GnRH increased the number of ovulating females compared with the use of CPE. The fertility of females injected with GnRH was also higher. It was found later that threshold doses of certain synthetic analogues of mammalian and salmon GnRH that would induce ovulation in female tench were significantly lower (Kouřil et al., 1989, 1991). Similar review studies of hormonally induced reproduction in female tench were published by Kouřil and Hamáčková (1996) and Kouřil (1998).

To compare and evaluate the effects of ovulation induction in female tench using GnRH_a and CPE, the results of biochemical examinations of blood plasma were also used.

The aims of the present study were:

- to induce ovulation and perform artificial propagation of female tench by repeated GnRH analogue administration in two consecutive spawning seasons
- to evaluate the effects of repeated administration of different hormonal preparations on weight gain and mortality of breeding fish, on success in achieving ovulation, artificial propagation and fecundity compared with females that were repeatedly given CPE
- to compare the values of selected blood plasma biochemical parameters between the two above groups of female tench
- to compare selected blood plasma biochemical parameters between stripped and not stripped tench females with ovulation induced by either GnRH_a or CPE

MATERIAL AND METHODS

The fish used in the experiments came from pond broodstocks of the Research Institute of Fish Culture and Hydrobiology in Vodňany. At the beginning of long-term monitoring, breeding tench were individually cryogenically marked. The fish were maintained in small experimental ponds 0.08 ha in size. In spring before spawning, the fish

were collected, males and females were separated and released to flow-through troughs. Before the first-year propagation, the fish were randomly divided into two groups of 19 females each. The mean weight of individual CPE-injected and GnRH_a-injected fish was 776.7 ± 19.63 and 816.7 ± 67.20 , respectively. In the latter group, ovulation was induced by the intramuscular administration of $10 \mu\text{g}/\text{kg}$ of synthetic superactive analogue of gonadotropin-releasing hormone / (D-Ala⁶) GnRHProNH₂Et/, in the former group by the administration of $2 \text{ mg}/\text{kg}$ dehydrated CPE. The average water temperature from administration to ovulation was 22.15°C in the first year and 22.25°C in the second year.

To assess the physiological conditions of production tench females in the present study, selected blood plasma biochemical indices divided into the following groups were used (Masopust, 2000):

- stress indices (cortisol, glucose);
- condition indices (total proteins – TP, triacyl glycerol – TGA);
- tests of permeability and integrity of cell membranes (alanine aminotransferase – ALT, aspartate aminotransferase – AST, creatine kinase – CK).

In the first year, blood samples of tench females ($N = 37$) were taken on 20 June about 0 to 6 hours after stripping. The fish were kept in troughs with flowing pond water. Water temperature was about 21°C . The next year, the tench were stripped on 16 June and blood sampling was performed on 17 June, i.e. about 24 hrs after stripping and around 48 hrs following the induction of ovulation. Blood samples were taken from 29 female tenches. At the time of blood sampling, water temperature was 22°C , and the fish were kept in troughs with flowing pond water. The fish were anesthetized with 2-phenoxyethanol before blood samples were taken (Hamáčková et al., 2004).

Blood (approximately 1 ml) was sampled from a. et v. caudalis per fish. Heparin sodium salt (50 i.u. per 1 ml blood) was used as anticoagulant (Svobodová et al., 1991). The radioimmunoanalytic method (Immunotech commercial kit) was used for the determination of blood plasma cortisol concentrations. Glucose concentrations, total proteins (TP), triacyl glycerol (TGA), transaminase (ALT, AST) and creatine kinase (CK) were determined using the automatic analyzer COBAS MIRA (F. Hoffmann, La-Roche and Co., Switzerland). Because the CK activity values were very high, blood plasma was diluted 5 to 10 times with physiological solution.

Table 1. Results of the first reproduction season

Preparation	Females		Relative weight of stripped eggs (%)	Latency period (°h)
	injected (N)	ovulated (%)		
GnRHa	17	63.2	6.19	674 ± 5
CPE	18	44.4	4.11	491 ± 34

Table 2. Results of the second reproduction season

Preparation	Females		Relative weight of stripped eggs (%)	Latency period (°h)
	injected (N)	ovulated (%)		
GnRHa	13	53.8	4.10	593 ± 5
CPE	16	37.5	3.40	531 ± 34

The results were statistically analyzed by means of one-way ANOVA.

RESULTS

In the first year of artificial stripping using GnRHa, 63.2% of female tench ovulated and the relative weight of stripped eggs was 6.19%. The mean absolute fecundity (with only stripped fish included) was 81 314 eggs per female. The relative fecundity (with only stripped fish included) was 96 528 eggs per kg. Ovulation in female tench treated with CPE was achieved in 44.4% of the cases and the relative weight of stripped eggs was 4.11%. The mean absolute fecundity in this group was 51 965 eggs per female, and the mean relative fertility was 68 062 eggs per kg. The latency period (between injection and ovulation) was 674 ± 5 °h and 491 ± 34 °h in fish injected with GnRHa and pituitary, respectively (Table 1).

Of the females injected with CPE and GnRH analogue in the first year, 84.2% and 72.2%, respectively, survived until the next year's spawning. The mean weight of females treated in the first year with CPE and GnRHa was 1 042.5 ± 150.4 and 998.8 ± 188.1, respectively. In the second spawning season, 37.5% fish injected with pituitary and 53.8% fish injected with GnRH analogue successfully ovulated and were stripped. The relative weight of stripped eggs, however, was lower by 4.1% and 3.4% after the administration of CPE and GnRHa, respectively. Both the relative and the absolute fecundity figures were also lower in both cases. In the second experimental year, the latency period was 531 ± 86 °h (fish injected with carp pituitary extract) and 593 ± 60 °h (fish injected with GnRHa) (Table 2).

Table 3 (year 1) and 4 (year 2) give the blood plasma biochemical profiles of studied female tench divided according to the method of ovulation induction into a group of (1) stripped female tench

Table 3. Comparison of blood plasma biochemical indices in stripped and not stripped tench females (year 1)

Index	GnRHa (stripped)	GnRHa (not stripped)	CPE (stripped)	CPE (not stripped)
	mean ± SD (n = 11)	mean ± SD (n = 6)	mean ± SD (n = 9)	mean ± SD (n = 9)
Fish body weight (g)	841.0 ± 126.67	774.8 ± 150.40	771.4 ± 121.35	779.2 ± 134.31
Glucose (mmol/dm ³)	8.80 ± 2.145 ^a	6.84 ± 2.299 ^{ab}	10.43 ± 3.831 ^a	5.93 ± 2.098 ^b
TP (g/dm ³)	36.8 ± 5.47 ^{ab}	42.17 ± 4.738 ^b	33.03 ± 5.440 ^a	34.92 ± 4.180 ^a
TGA (mmol/dm ³)	1.54 ± 0.184 ^a	1.53 ± 0.641 ^a	1.27 ± 0.252 ^{ab}	1.08 ± 0.296 ^b
ALT (µkat/dm ³)	0.25 ± 0.125	0.23 ± 0.132	0.19 ± 0.091	0.16 ± 0.059
AST (µkat/dm ³)	4.09 ± 2.237	5.15 ± 2.418	4.76 ± 0.964	4.63 ± 1.741
CK (µkat/dm ³)	87.7 ± 60.05	134.3 ± 88.11	102.2 ± 39.94	93.8 ± 54.85

The values with superscript^{a,b} express significant differences between the groups compared

with ovulation induced by GnRH α , and (2) stripped female tench with ovulation induced by CPE. It follows from the results that there were no differences in the biochemical indices of stress, condition or values characterizing the permeability and integrity of cell membranes between the groups of stripped female tench where GnRH α or CPE was used to induce ovulation. The same results were obtained in the first and in the second year of the experiment.

Another objective of the study was to compare the biochemical parameters of blood plasma between four groups of tench females, stripped and not stripped with ovulation induced by either GnRH α or CPE. The results of analyses and comparisons of these groups of female tench are shown in Table 3 (year 1) and 4 (year 2). The values of condition indices (TP, TGA) and of the cell membrane characteristics (ALT, AST, CK) in stripped and not stripped female tench following the application of either GnRH α or CPE immediately after stripping (year 1) and about 24 hrs after stripping (year 2) were practically comparable without any significant differences. Different values were, however, found in stress indices, i.e. in glucose and cortisol concentrations. In year 1 of female tench monitoring, higher blood plasma glucose levels were recorded immediately after stripping in stripped tench compared with not stripped females; in tench that had been administered CPE the difference was significant ($P < 0.01$) compared with fish treated with GnRH α . In year 2 when blood samples were collected about 24 hours after stripping, on the other hand, higher glucose and cortisol blood plasma concentrations were found in all not stripped female tench. The differences in glucose concentrations were significant in both the group of female tench treated with GnRH α ($P < 0.01$) and that treated with CPE ($P < 0.05$).

DISCUSSION

It follows from the results that the artificial spawning of female tench with repeated hormonally induced ovulation may significantly increase mortality in subsequent years. The assumption of higher mortality among females injected with CPE compared with females injected with GnRH analogue, however, was not confirmed. In fact, the results suggest that the opposite is true. Lower weight gains among tench females repeatedly administered

the GnRH analogue compared with females that were administered the carp pituitary were found. In both investigated periods, the number of successfully spawning females was about 30 per cent higher in the group where ovulation had been induced by the GnRH analogue than in the group of females that had been administered CPE. The relative weight of stripped eggs (related to the weight of female tench before stripping) in two consecutive spawning seasons in females with artificially induced stripping showed a markedly decreasing trend. In the investigated period, the relative weight of stripped eggs was 1/4–1/2 lower in the group of females administered CPE than in the group of females administered the GnRH analogue.

Reproduction in fish is one of the factors that significantly influence the internal environment of the organism. For that reason, much attention is paid to the study of haematological and biochemical indices in the reproductive period. Changes in haematological indices in tench during their reproductive season were described by Einszporn-Orecka (1970) and Svobodová et al. (1978). Einszporn-Orecka (1970) reported a decrease in the values of haematological indices (erythrocyte count Er, haemoglobin count Hb, hematocrit value PCV) during the spawning season in breeding tench in lakes, and reported a marked decrease in the values in females. A similar character of changes in haematological indices (Er, Hb, PCV) in female and male tench after artificial propagation was described by Svobodová et al. (1978).

The present study reports the results of biochemical indices evaluating stress, condition and the characteristics of cell membranes in female tench after artificial propagation. No differences in biochemical profiles in the two years of investigation were found between the two groups of stripped female tench whose ovulation had been induced by GnRH α and CPE, respectively. In their study in female wels, Svobodová et al. (1997) found no differences in either haematological (Er, Hb, PCV, MCV, MCH, MCHC, leukocytes) or biochemical indices (TP, glucose, cortisol) when they used different hormonal substances (GnRH and CPE) to induce ovulation.

When ovulation was induced by GnRH α or CPE, only some females spawned while others failed to spawn. No significant differences in biochemical indices characterizing the condition and characteristics of cell membranes were found between the stripped and not stripped groups of females. Different values were recorded in stress indices,

Table 4. Comparison of blood plasma biochemical indices in stripped and not stripped tench females (year 2)

Index	GnRHa (stripped)	GnRHa (not stripped)	CPE (stripped)	CPE (not stripped)
	mean \pm SD ($n = 8$)	mean \pm SD ($n = 5$)	mean \pm SD ($n = 6$)	mean \pm SD ($n = 10$)
Fish body weight (g)	985.4 \pm 223.10	1 020.2 \pm 134.57	1 066.2 \pm 109.70	1 056.8 \pm 158.44
Cortisol (mmol/dm ³)	519.9 \pm 172.67	622.6 \pm 127.69	334.8 \pm 228.28	517.3 \pm 232.74
Glucose (mmol/dm ³)	9.09 \pm 1.24 ^a	12.67 \pm 2.751 ^b	7.10 \pm 1.992 ^a	11.00 \pm 3.969 ^b
TP (g/dm ³)	38.7 \pm 5.92	33.6 \pm 4.15	37.4 \pm 4.14	35.0 \pm 5.29
Tcg (mmol/dm ³)	2.45 \pm 0.679	2.07 \pm 0.385	2.16 \pm 0.531	2.11 \pm 0.315
ALT (μ kat/dm ³)	0.15 \pm 0.075	0.15 \pm 0.065	0.16 \pm 0.098	0.19 \pm 0.058
AST (μ kat/dm ³)	6.90 \pm 1.154	6.39 \pm 1.496	7.40 \pm 2.073	6.83 \pm 1.657
CK (μ kat/dm ³)	205.0 \pm 162.98	165.7 \pm 27.41	201.3 \pm 188.77	236.3 \pm 133.93

The values with superscript^{a,b} express significant differences between the groups compared

specifically in glucose concentrations. A significant increase ($P < 0.01$) in glucose concentrations was observed in females immediately after stripping. This was a persisting response to the physiological and handling stress from the stripping period. Similar values were reported by Jeney and Jeney (1992) and Svobodová et al. (1997), Svoboda et al. (2001) in the blood plasma of female carp, wels and tench immediately after artificial spawning. In not stripped female tench, significantly higher ($P < 0.01$) concentrations of glucose were found about 48 hours after the ovulation induction compared with stripped females about 24 hours after stripping. This was probably in response to the stress produced by the inability to release eggs, by the eggs remaining in the body cavity, etc. The stress also manifested itself by a slight, insignificant decrease in total protein and triacyl glycerol concentrations in the blood plasma of unspawned females (Table 4). It follows from the above that both artificial stripping and the inability to release eggs are important stress factors for female tench. This conclusion underlines the importance of welfare.

Hardly any investigated parameter showed differences between the groups of females whose ovulation had been induced by GnRHa or by CPE. The use of GnRHa for the induction of ovulation in artificially spawned tench is preferable because it seems to produce better results in absolute and relative fertility.

REFERENCES

Cooperative Team (1977): Cooperative Team for hormonal application in pisciculture. A highly effective ovu-

lating agent for fish reproduction practical application of LH-RH analogue for the induction of spawning of farm fishes. *Sci. Sin.*, 20, 469–474.

Donaldson E.M., Hubte C.A., Dye H.M. (1981): Induced ovulation in Pacific salmon using LHRH analogue and salmon gonadotropin. In: 9th Symp. Comp. Endocrinol., Hong Kong, 137 pp.

Einszporn-Orecka T. (1970): Quantitative changes in the circulating blood of tench (*Tinca tinca* L.) in the annual cycle. *Pol. Arch. Hydrobiol.*, 17, 435–444.

Hamáčková J., Lepičová A., Kozák P., Stupka Z., Kouřil J., Lepič P. (2004): The efficacy of various anaesthetics in tench (*Tinca tinca* L.) related to water temperature. *Vet. Med.–Czech.*, 12, 467–472.

Jeney Z., Jeney G. (1992): Primary and secondary stress responses of common carp (*Cyprinus carpio* L.) caused by artificial propagation. In: Adámek Z., Flajšhans M. (eds.): Proc. Conf. Fish Reproduction, Vodňany, 27–30.

Kouřil J. (1987): Induced ovulation of the female tench (*Tinca tinca* L.): Effect of the number and levels of pituitary injections on the results of stripping. *Práce VÚRH Vodňany*, 16, 53–61. (in Czech)

Kouřil J. (1998): Hormonally induced spawning of tench *Tinca tinca* (L.) females (A review). *Pol. Arch. Hydrobiol.*, 45, 421–433.

Kouřil J., Barth T. (1981): The achievement of egg ovulation in artificial spawning of tench (*Tinca tinca* L.) using LH-RH. *Bul. VÚRH Vodňany*, 17, 13–18. (in Czech with English summary)

Kouřil J., Hamáčková J. (1996): Hormonally induced artificial spawning of tench (*Tinca tinca* L.) females using synthetic GnRH analogues. In: Flajšhans M. (ed.): Proc. Scientific Papers the 75th Anniversary of Foundation of the Research Institute of Fish Culture and Hydrobiology. Vodňany RIFCH USB, 49–60.

- Kouřil J., Barth T., Hamáčková J., Flegel M. (1986): Induced ovulation in tench (*Tinca tinca* L.) by various LH-RH synthetic analogues: effect of site of administration and temperature. *Aquaculture*, 54, 37–44.
- Kouřil J., Mikodina E.V., Glubokov A.I., Hamáčková J., Barth T., Flegel M., Charvátová J. (1989): The utilization of /Glu(NH-Ad⁶), Trp⁷, Leu⁸/ Gn-RH for the induction of tench (*Tinca tinca* L.) females ovulation. *Bul. VÚRH Vodňany*, 25, 8–13. (in Czech with English summary)
- Kouřil J., Barth T., Hamáčková J., Mikodina E.V., Glubokov A.I., Flegel M. (1991): The comparison of effects hormonal induction of ovulation in tench (*Tinca tinca* L.) in artificial propagation. In: *Proc. Int. Symp. Reproductive Biology in aquaculture*, Taipei, Taiwan, 54 pp.
- Masopust J. (2000): *Clinical Biochemistry*. Karolinum, Prague. 832 pp. (in Czech)
- Pokorný J. (1974): Spawning of the tench and culture of the fry. *Rybářství*, 12, 268–270. (in Czech)
- Svoboda M., Kouřil J., Hamáčková J., Kaláb P., Savina L., Svobodová Z., Vykusová B. (2001): Biochemical profile of blood plasma of tench (*Tinca tinca* L.) during pre- and postspawning period. *Acta Vet. Brno*, 70, 259–268.
- Svobodová Z., Kouřil J., Hamáčková J. (1978): The values of some haematological indices in parent tench (*Tinca tinca* L.). *Živoč. Výr.*, 23, 825–833. (in Czech with English summary)
- Svobodová Z., Pravda D., Paláčková J. (1991): Unified methods of haematological examination of fish. *Research Institute of Fish Culture and Hydrobiology, Vodňany*, 31 pp. (in Czech)
- Svobodová Z., Kolářová J., Kouřil J., Hamáčková J., Vykusová B., Kaláb P. (1997): Haematological investigations in *Silurus glanis* L. females during pre- and postspawning period. *Pol. Arch. Hydrobiol.*, 44, 67–81.

Received: 2006–05–17

Accepted after corrections: 2007–04–02

Corresponding Author

Assoc. Prof. Ing. Jan Kouřil, Ph.D., University of South Bohemia České Budějovice, Research Institute of Fish Culture and Hydrobiology, Zátíší 728/II, 389 25 Vodňany, Czech Republic
Tel. +420 383 382 402, e-mail: kouril@vurh.jcu.cz
