

## 海南近岸海域夏初角毛藻的种类多样性及其水平分布

庞勇<sup>1</sup>, 聂瑞<sup>2</sup>, 王道儒<sup>1</sup>, 黄强<sup>3</sup>

1. 海南省海洋与渔业科学院, 海南 海口 570125; 2. 海南省海洋监测预报中心, 海南 海口 570206; 3. 山东恒诚检测科技有限公司, 山东 烟台 261400

**Species diversity and horizontal distribution of Chaetoceros in coastal waters of Hainan Island during early summer****PANG Yong<sup>1</sup>, NIE Rui<sup>2</sup>, WANG Daoru<sup>1</sup>, HUANG Qiang<sup>3</sup>**

1. Hainan Academy of Oceanographic and Fishery Sciences, Haikou 570125, China; 2. Marine Monitoring and Forecasting Center of Hainan Province, Haikou 570206, China; 3. Shandong Hengcheng Testing Technology Co. Ltd., Yantai 261400, China

[摘要](#)[图/表](#)[参考文献\(0\)](#)[相关文章 \(1\)](#)[全文: PDF \(2248 KB\)](#)[HTML \(0\)](#)[输出: BibTeX | EndNote \(RIS\)](#)**摘要**

根据2011年5月28日至6月3日在海南近岸海域采集的浮游植物样品的鉴定结果, 分析了角毛藻(*Chaetoceros*)的种类多样性、优势种构成、细胞数量和水平分布。结果显示, 海南近岸海域共发现角毛藻28种(含变种、变型), 其中西北部海域25种, 东部海域22种, 南部海域21种; 暹罗角毛藻(*C.iamense*)、拟旋链角毛藻(*C.pseudocurvisetus*)、并基角毛藻(*C.decipiens*)和劳氏角毛藻(*C.lorenzianus*)为主要优势种, 另外, 桥联角毛藻(*C.anastomosans*)、嘴状角毛藻(*C.rostratus*)、密链角毛藻(*C.densus*)、窄隙角毛藻(*C.affinis*)和短叉角毛藻(*C.messanensis*)的数量也较多。东部海域角毛藻细胞数量最高, 平均为(162.70±415.16)×10<sup>4</sup>个·m<sup>-3</sup>; 西北部海域角毛藻数量略低于东部海域, 平均为(102.81±206.26)×10<sup>4</sup>个·m<sup>-3</sup>; 南部海域角毛藻数量最低, 平均为(70.43±142.09)×10<sup>4</sup>个·m<sup>-3</sup>, 角毛藻细胞数量平面分布呈现由近岸向外海递减的趋势, 与环境因子之间不存在显著相关关系。

**关键词:** 海南近岸, 角毛藻, 种类多样性, 水平分布**Abstract :**

Based on a survey on the species diversity, dominant species composition, cell abundance and horizontal distribution of *Chaetoceros* in the coastal waters of Hainan conducted during May 28 and June 3 in 2011, we identified 28 species of *Chaetoceros*, among which 25 species were found in the northwestern waters, 22 in the eastern waters and 21 in the southern waters. The dominant species in these coastal waters were *C.iamense*, *C.pseudocurvisetus*, *C.decipiens* and *C.lorenzianus*, while *C.anastomosans*, *C.rostratus*, *C.densus*, *C.affinis* and *C.messanensis* were also abundant in these coastal waters of Hainan. Eastern waters had the highest cell abundance with an average number of (162.70±415.16)×10<sup>4</sup> ind·m<sup>-3</sup>, while northwestern waters was slightly lower in cell abundance with an average number of (102.81±206.26)×10<sup>4</sup> ind·m<sup>-3</sup>, and southern waters had a minimum value of (70.43±142.09)×10<sup>4</sup> ind·m<sup>-3</sup>. Horizontal distribution of *Chaetoceros* in these three waters had a declining trend from onshore to offshore. There was no significant correlation between *Chaetoceros* abundance and environmental factors.

**Key words:** coastal waters of Hainan    *Chaetoceros*    species diversity    horizontal distribution**收稿日期:** 2014-10-15    **修回日期:** 2015-02-05    **出版日期:** 2015-12-05**PACS:** Q 145

Q 178.53

**基金资助:**

海岸带主体功能区划分技术框架研究示范项目(200905005-02); 海洋公益性行业科研专项经费项目(201005012-6)

**作者简介:** 庞勇(1983-), 男, 硕士, 助理研究员, 从事海洋生态环境研究。E-mail: yongpang520@163.com**引用本文:**

庞勇, 聂瑞, 王道儒, 黄强. 海南近岸海域夏初角毛藻的种类多样性及其水平分布[J]. 南方水产科学, 2015, 11(6): 9-18. PANG Yong, NIE Rui, WANG Daoru, HUANG Qiang. Species diversity and horizontal distribution of Chaetoceros in coastal waters of Hainan Island during early summer. South China Fisheries Science, 2015, 11(6): 9-18.

**链接本文:**

<http://www.schinafish.cn/CN/10.3969/j.issn.2095-0780.2015.06.002> 或 <http://www.schinafish.cn/CN/Y2015/V11/I6/9>



