



 **Current Issue**


 **Browse Issues**


 **Search**



 **About this Journal**

 **Instruction to Authors**

 **Online Submission**

 **Subscription**

 **Contact Us**



 **RSS Feed**

## Acta Medica Iranica

2009;47(4) : 204-211

### RESISTANCE OF ANOPHELES SACHAROVII FAVRE TO DDT

AV. Manouchehri, A.Zaini, E. Javadian, E. Saebi

#### Abstract:

A.sacharovi, which is one of the main malaria vectors of the Palearctic region, is resistant to DDT in Greece (1949), Lebanon (1954), Turkey (1959), Syria and the U.S.S.R. (1972).The first record of A. sacharovi resistance to DDT in Iran was given in 1959 from the Kazeroon area, followed by Izeh (1969) and Meshkin-Shahr (1970).This study (1973) urgently advises speedy measures to eliminate the present foci of DDT –resistant A.sacharovi, to distinguish the checking of the susceptibility level of A. sacharovi in other areas of its distribution in Iran.

#### Keywords:

[Malaria vectors](#) , [Palearctic region](#)

TUMS ID: 2008

Full Text HTML  Full Text PDF  280 KB

top ▲

[Home](#) - [About](#) - [Contact Us](#)

TUMS E. Journals 2004-2009  
Central Library & Documents Center  
Tehran University of Medical Sciences

Best view with Internet Explorer 6 or Later at 1024\*768 Resolutions