



 **Current Issue**


 **Browse Issues**

 **Search**



 **About this Journal**


 **Instruction to Authors**

 **Online Submission**

 **Subscription**

 **Contact Us**



 **RSS Feed**

## Acta Medica Iranica

2009;47(4) : 47-54

### HUMAN DICROCOELIASIS 2 CASE REPORTS FROM IRON

A.Sohrabi

#### Abstract:

Dicrocoelium dendriticum is 10 by 2 mm. trematodes which live in the liver of the cattle, goats and sheep but it occasionally infects the human being. During the study of intestinal infections in the Caspian Sea region 2 cases of Dicrocoeliasis were found. The method for diagnosis was Formol-Ether concentration technique for stool examination. These two patients had eosinophilia and vague abdominal pains and digestive disorders. The following points are worthwhile to remind for the patients in the Caspian Sea region. 1. The Formol-Ether technique of stool examination is the best method for diagnosis of trematodes ova. 2. Those patients who come from the Caspian Sea region and have vague abdominal pain and eosinophilia should have a stool examination by the Formol-Ether technique. 3. In the differential diagnosis of ictrus and liver diseases one should remember the distomatosis.

TUMS ID: 1623

Full Text HTML  Full Text PDF  313 KB

top ▲

[Home](#) - [About](#) - [Contact Us](#)

TUMS E. Journals 2004-2009  
Central Library & Documents Center  
Tehran University of Medical Sciences

Best view with Internet Explorer 6 or Later at 1024\*768 Resolutions