

CASE REPORT

[Previous Article](#) [ToC](#) [Next Article](#)

Year : 2007 | Volume : 41 | Issue : 4 | Page : 401-403

Tarlov cyst: Case report and review of literature

Bhagwat Prashad, Anil K Jain, Ish K Dhammi

University College of Medical Sciences and Guru Teg Bahadur Hospital, Shahadara, Delhi, India

Click [here](#) for **correspondence address** and email

Download Article (pdf)
 Email Article
 Print Article
 Read / Write a Comment
 Citation Manager

Abstract

We describe a case of sacral perineural cyst presenting with complaints of low back pain with neurological claudication. The patient was treated by laminectomy and excision of the cyst. Tarlov cysts (sacral perineural cysts) are nerve root cysts found most commonly in the sacral roots, arising between the covering layer of the perineurium and the endoneurium near the dorsal root ganglion. The incidence of Tarlov cysts is 5% and most of them are asymptomatic, usually detected as incidental findings on MRI. Symptomatic Tarlov cysts are extremely rare, commonly presenting as sacral or lumbar pain syndromes, sciatica or rarely as cauda equina syndrome. Tarlov cysts should be considered in the differential diagnosis of patients presenting with these complaints.

Keywords: Low back pain, sacral perineural cyst, sciatica, tarlov cyst

How to cite this article:

Prashad B, Jain AK, Dhammi IK. Tarlov cyst: Case report and review of literature. Indian J Orthop 2007;41:401-3

How to cite this URL:

Prashad B, Jain AK, Dhammi IK. Tarlov cyst: Case report and review of literature. Indian J Orthop [serial online] 2007 [cited 2008 Nov 20];41:401-3. Available from: <http://www.ijonline.com/text.asp?2007/41/4/401/37007>

Tarlov cysts were first described in 1938 as an incidental finding at autopsy. ^[1] Tarlov described a case of symptomatic perineural cyst and recommended its removal. Since then a few cases have been reported in the literature. ^{[2],[3],[4]}

Paulsen reported the incidence of Tarlov cysts as 4.6% in back pain patients (n=500). Only 1% of back pain patients (n=500) were symptomatic. ^[4] The patient may present as low back pain, sciatica, coccydynia or cauda equina syndrome. The cysts are usually diagnosed on MRI, which reveals the lesion arising from the sacral nerve root near the dorsal root ganglion. ^[5]

Search

SEARCH

Similar in PUBMED

Search Pubmed for

- Prashad B
- Jain AK
- Dhammi IK

Email Alert *

Add to My List *

* Registration required (free)

Abstract

Case Report

Discussion

References

Article Figures

Article Access Statistics

Viewed	3499
Printed	34
Emailed	3
PDF Downloaded	126
Comments	[Add]



Figures and Tables



Tarlov advised extensive surgery with sacral laminectomy and excision of the cyst along with the nerve root. [6] Paulsen reported CT-guided percutaneous aspiration of these perineural cysts for relief of sciatica. [4] Recently, microsurgical excision of the cyst has been advocated, combined with duraplasty or plication of the cyst wall. [7]

Recommend
this journal
for your library



We report a case of symptomatic Tarlov cyst presenting as back pain, to increase the awareness of this rare entity in the orthopedic community.

Case Report

A 29-year-old female presented with right thigh pain off and on for nine months. The pain was not associated with specific time, posture or activity and it used to get relieved by non steroidal antiinflammatory drugs (NSAID). Clinical examination at this stage did not reveal any findings at spine, bilateral hips and left thigh.

For last three months, the intensity and duration of pain had increased, which was now unrelieved by taking NSAID. The pain had progressed to the lower back and bilateral upper thigh up to the ankle. The pain was aggravated by activity and prolonged standing and was more bothersome in the evening. She used to get up in the middle of the night with pain. Later the patient started having rest pain as well. Examination showed no spinal tenderness. Straight leg raising was 50° on the right side and normal on the left side. There was mild blunting of sensations along the S1 and S2 dermatome on the right side, no motor deficit in both lower limbs.

X-ray of the lumbosacral spine did not reveal any abnormality [Figure - 1]. The MRI of the spine revealed fluid-filled cystic lesion, arising from the second sacral nerve root on the right side and measuring 2cm in diameter [Figure - 2].

The patient was taken for sacral laminectomy, excision of the cyst and plication of the cyst wall, while retaining the nerve root [Figure - 3]. Histopathological examination of the cyst wall showed nerve cells, which confirmed the diagnosis of Tarlov cyst.

Patient appreciated relief of pain immediately after the surgery. Postoperative period was uneventful and the patient made prompt recovery. On nine months followup, the patient had no pain in lower limbs and back. The patient is back at her job and is asymptomatic. Postoperative MRI taken at nine months [Figure - 4] did not show any evidence of recurrence of the cyst.

Discussion

Tarlov cysts are rare causes of low back pain. They are more common in females. [4],[7] Clinical presentation of Tarlov cysts is variable. The cysts may cause local and/or radicular pain. The dominant syndrome is referable to the caudal nerve roots, either sciatica, sacral or buttocks pain, vaginal or penile paraesthesia or sensory changes over the buttocks, perineal area and lower extremity. Depending on their location, size and relationship to the nerve roots, they may cause sensory disturbances or motor deficits to the point of bladder dysfunction. Tenderness on firm pressure over the sacrum may be present. Commonly, the symptomatology is intermittent at its onset and is most frequently exacerbated by standing, walking and coughing. Bed rest alleviates the discomfort. [8]

Plain X-rays are usually normal. However, they may reveal characteristic bone erosion of the spinal

canal or anterior or posterior neural foramina. [9] A CT scan can demonstrate cystic masses isodense with CSF located at the foramina. Bony changes may also be present. [10] An MRI gives a much better soft tissue contrast and is currently the investigation of choice for perineural cysts. The cysts demonstrate low signal on T-1 weighted images and high signal on T-2 weighted images, similar to CSF. [5] Myelography showing the filling of the meningocele sac 1h after injection of contrast medium is highly suggestive of a perineural cyst. [11]

Microscopic features of the cyst were described by Tarlov. The early stage in cyst formation is that of a space between the arachnoid which covers the root or the perineurium and the outer layer of the pial cover of the root or the endoneurium. It usually begins in one portion of the circumference of the perineural space, the larger cysts compressing the nerve root to one side. The cyst occupies the posterior root abutting the proximal portion of the dorsal ganglion. Its main part is bordered by reticulum or by nerve fibers. [1]

The pathogenesis of perineural cysts is uncertain. Tarlov felt that hemorrhage into the subarachnoid space caused accumulations of red cells which impeded the drainage of the veins in the perineurium and epineurium, leading to rupture with subsequent cyst formation. Four out of the seven patients in Tarlov's 1970 article had a history of trauma. [8] Schreiber and Haddad also supported this posttraumatic cause of cyst formation. [12] Because many of the patients with perineural cyst in their series did not have histories of trauma, Fortuna *et al* . believed that the perineural cysts were congenital, caused by arachnoidal proliferations within the root sleeve. [13]

There is no consensus on a single method of treatment. Various methods have been advocated. Tarlov advised that symptomatic, single perineural cysts should be completely excised together with the posterior root and ganglion from which they arise. [8] Paulsen reported CT-guided percutaneous aspiration of these perineural cysts in two patients for the relief of sciatica caused by compression. [4] According to Caspar microsurgical excision of the cyst combined with duraplasty or plication of the cyst wall is an effective and safe treatment of symptomatic sacral cysts. The parent nerve root is always left intact. [7]

Tarlov cysts are a documented cause of sacral radiculopathy and other radicular pain syndromes. They must be considered in the differential diagnosis of patients presenting with these clinical presentations and appropriately treated by cyst excision.

References

1. Tarlov IM. Perineural cysts of the spinal nerve roots. *Arch Neural Psychiatry* 1938;40:1067-74. [↑](#)
2. Chaiyabud P, Suwanpratheep K. Symptomatic Tarlov cyst: Report and review. *J Med Assoc Thai* 2006;89:1047-50. [↑](#) [PUBMED]
3. Acosta FL Jr, Quinones-Hinojosa A, Schmidt MH, Weinstein PR. Diagnosis and management of sacral Tarlov cysts. Case report and review of the literature. *Neurosurg Focus* 2003;15:E15. [↑](#)
4. Paulsen RD, Call GA, Murtagh FR. Prevalence and percutaneous drainage of cysts of the sacral nerve root sheath (Tarlov cysts). *AJNR Am J Neuroradiol* 1994;15:293-9. [↑](#) [PUBMED]
5. Rodziewicz GS, Kaufman B, Spetzler RF. Diagnosis of sacral perineural cysts by nuclear magnetic resonance. *Surg Neurol* 1984;22:50-2. [↑](#) [PUBMED]
6. Tarlov IM. Cysts (perineurial) of the sacral roots. *J Am Med Assoc* 1948;138:740-4. [↑](#)
7. Caspar W, Papavero L, Nabhan A, Loew C, Ahlhelm F. Microsurgical excision of symptomatic sacral perineurial cysts: A study of 15 cases. *Surg Neurol* 2003;59:101-6. [↑](#) [PUBMED] [FULLTEXT]
8. Tarlov IM. Spinal perineurial and meningeal cysts. *J Neural Neurosurg Psychiatry* 1970;33:833-43. [↑](#)
9. Taveras JM, Wood EH. *Diagnostic neuroradiology*. 2nd ed. Vol 2. Williams and Wilkins: Baltimore; 1976. p.1139-45. [↑](#)
10. Tabas JH, Deeb ZL. Diagnosis of sacral perineural cysts by computed tomography. *J Comput Tomogr* 1986;10:255-9. [↑](#)

11. Nishiura I, Koyama T, Handa J. Intracranial perineurial cyst. *Surg Neurol* 1985;23:265-9. [↑](#)
 12. Schreiber F, Haddad B. Lumbar and sacral cysts causing pain. *J Neurosurg* 1951;8:504-9. [↑](#)
 13. Fortuna A, La Torre E, Ciappetta P. Arachnoid diverticula: A unitary approach to spinal cysts communicating with the subarachnoid space. *Acta Neurochir (Wien)* 1977;39:259-68. [↑](#)
-

▲

Correspondence Address:

Bhagwat Prashad
K-1/133, First Floor, Chitaranjan Park, New Delhi-110 019
India

 [Login to access the email ID](#)

Source of Support: None, **Conflict of Interest:** None

 **Figures**

[Figure - 1], [Figure - 2], [Figure - 3], [Figure - 4]

▲

    

[◀ Previous Article](#) [Next Article ▶](#)

[Contact us](#) | [Sitemap](#) | [Advertise](#) | [What's New](#) | [Feedback](#) | [Copyright and Disclaimer](#)

© 2006 - Indian Journal of Orthopaedics | A journal by Medknow

Online since 9th November, 2006