

 [Current Issue](#) [Browse Issues](#) [Search](#) [About this Journal](#) [Instruction to Authors](#) [Online Submission](#) [Subscription](#) [Contact Us](#) [RSS Feed](#)

Acta Medica Iranica

2009;47(4) : 107-110

Original Article

Abdominal Incisional Endometriosis Following Cesarean Section: Report of 2 Cases

Azam Azargoon, M.D.;1 Nahid Azargoon, M.D.;2

1 Department of Gynecology and Obstetrics, Semnan University of Medical Sciences, Semnan, Iran

2 Shafa Hospital, Semnan, Iran

Corresponding Author:

Azam Azargoon, Amir- Al- Momenin Hospital, Madar square, Semnan, Iran.

P.O. Box: 35197-34731

Tel: 0098-2314460088 Fax: 0098-2314461580

E-mail: azarmona2003@yahoo.com

Received: October 16,2007

Accept : December 8,2007

Abstract:

Endometriosis is described as the presence of functioning endometrial tissue (glands and stroma) outside the uterine cavity. The most common location is within the pelvis. However, extra pelvic endometriosis is a fairly uncommon disorder and difficult to diagnose. It can sometimes occur in a surgical scar. Scar endometriosis is a rare condition and difficult to diagnose. It mostly follows obstetrical and gynecological surgeries. It presents as a painful, slowly growing mass in or near a surgical scar. We report two cases of abdominal wall endometriosis following hysterotomy and cesarean section. Consequently the pathogenesis, diagnosis and treatment are discussed.

Keywords:

[Abdominal wall](#) , [Cesarean scar](#) , [Endometriosis](#)

TUMS ID: 12080

Full Text HTML  Full Text PDF  888 KB

top ▲

[Home](#) - [About](#) - [Contact Us](#)

TUMS E. Journals 2004-2009
Central Library & Documents Center
Tehran University of Medical Sciences

Best view with Internet Explorer 6 or Later at 1024*768 Resolutions