



- Current Issue
- Browse Issues
- Search
- About this Journal
- Instruction to Authors
- Online Submission
- Subscription
- Contact Us
- RSS Feed

## Acta Medica Iranica

2009;47(4) : 61-70

### THE CAUSATIVE AGENTS OF TINEA CRURIS

N. Moghadami, M.Emami

#### Abstract:

Within a period of 8 months from February to October 1986 345 Patients with suspected clinical mycotic lesions in the region of groin were referred to the Medical Mycology Department of the School of Public Health of Tehran Medical sciences University. Out of 345 Patients, 111 (9652%) Tinea cruris, 2 cases (1.74%); Tinea cruris associated with candidiasis, 1 case (0/87%);Tinea cruris associated with Erythrasma, 1 case (0/87%); Tinea cruris associated with Tinea versicolor were Observed. Agents which were isolated contained, E. floccosum 399 cases (11.11%) new species which are intermediate between T.mentagrophytes and T. rubrun 4 cases (4.94%), M.canis 3 cases (3.71%).T.violaceum 1 case (1.23%) and 1 case (1.23%) of Chrysosporim.

#### Keywords:

Epidermophyton floccosun . Trichophyton mentagrophytes

TUMS ID: 1559

Full Text HTML Full Text PDF 549 kB

top ▲

[Home](#) - [About](#) - [Contact Us](#)

TUMS E. Journals 2004-2009  
[Central Library & Documents Center](#)  
[Tehran University of Medical Sciences](#)

Best view with Internet Explorer 6 or Later at 1024\*768 Resolutions