



Turkish Journal of Medical Sciences

Turkish Journal

of

Medical Sciences

 [Keywords](#)
 [Authors](#)



medsci@tubitak.gov.tr

[Scientific Journals Home Page](#)

Are Inflammatory Cytokines Potential Mediators of Phenytoin-induced Gingival Overgrowth?

Nurcan BUDUNELI¹ Gül ATILLA¹ Necil KÜTÜKÇÜLER² ¹Department of Periodontology, Faculty of Dentistry, ²Department of Paediatrics, Faculty of Medicine, Ege University, İzmir - TURKEY Abstract : The levels of interleukin-1beta, tumour necrosis factor-alpha, and interleukin 6 in gingival crevicular fluid of phenytoin-treated patients (from sites exhibiting overgrowth and sites not exhibiting overgrowth) and control subjects were investigated. Six epileptic patients exhibiting severe phenytoin-induced gingival overgrowth, 12 patients with chronic gingivitis and 11 subjects with clinically healthy periodontium were included in the study. The cytokine levels were analysed by enzyme-linked immunosorbent assay and expressed as ng/2 sites. Wilcoxon signed-rank test, Kruskal-Wallis, Mann-Whitney U test and Spearman rank correlation were used for statistical analysis of the data. Overgrown sites showed significantly higher levels of interleukin-1beta compared to the healthy and gingivitis sites (p=0.0009, p=0.049 respectively). However, the difference was not statistically significant between overgrown and non-overgrown sites in epileptic patients (p=0.1159). Tumour necrosis factor-alpha values did not reveal any significant difference among the study groups. For interleukin 6, the highest value was found in the gingivitis sites, followed by overgrown, non-overgrown, and healthy sites, in descending order. No correlation was found between gingival crevicular fluid cytokine levels and papilla bleeding index values in phenytoin-treated patients. The results suggest that these cytokines play an indirect role in the complex mechanism of phenytoin-induced gingival overgrowth.

Key Words: phenytoin, interleukin-1beta, tumour necrosis factor-alpha, interleukin-6, gingival crevicular fluid

Turk J Med Sci 2001; **31**(1): 41-46.

Full text: [pdf](#)

Other articles published in the same issue: [Turk J Med Sci,vol.31,iss.1.](#)