JUSTAGE	My J-STAGE Sign in
The Bulletin of Tokyo Dental College	Published by Tokyo Dental College, Japan
Available Issues Japanese	>> <u>Publisher Site</u>
Author: ADVANCED	Volume Page
Keyword: Search	Go
Add to Favorite / Citation Add to Articles Alerts Publications Add to Favorite Favorite Alerts Publications Alerts HELP	

<u>TOP</u> > <u>Available Issues</u> > <u>Table of Contents</u> > Abstract

The Bulletin of Tokyo Dental College

Vol. 50 (2009), No. 3 :141-147

PRINT ISSN: 0040-8891

[PDF (313K)] [References]

## Six-year Follow-up in Skeletal Class III Patient Aged over 40 Receiving Orthognathic Surgery and Autotransplantation: A Case Report

Etsuko Motegi<sup>1)</sup>, Yumi Takane<sup>2)</sup>, Eri Tokunaga<sup>3)</sup>, Kenji Sueishi<sup>1)</sup>, Nobuo Takano<sup>4)</sup>, Takahiko Shibahara<sup>4)</sup> and Chikara Saito<sup>5)</sup>

1) Department of Orthodontics, Tokyo Dental College

2) Division of Dentistry, Takane Hospital

3) Division of Orthodontics, Department of Oral Health Clinical Science, Tokyo Dental College

4) Department of Oral and Maxillofacial Surgery, Tokyo Dental College

5) Division of Reconstructive Surgery for Oral and Maxillofacial Region, Department

of Tissue Regeneration and Reconstruction, Course for Oral Life Science, Niigata University Graduate School of Medical and Dental Sciences

> (Received April 30, 2009) (Accepted June 12, 2009)

**Abstract:** This paper describes the post-operative course of care in a patient requiring orthognathic surgery for skeletal mandibular protrusion in whom autotransplantation of a third molar was performed. A lower third molar that had to be removed for sagittal split ramus osteotomy (SSRO) was transplanted to replace the missing right second molar during pre-surgical orthodontic treatment, contributing to post-treatment occlusal stability. A 44-year-old woman presented with mandibular protrusion. The upper left second molar was congenitally missing and the lower right second molar had been extracted. She was diagnosed as having skeletal mandibular protrusion with excess vertical growth of the mandible and anterior open bite. Correction of the skeletal problem required orthognathic surgery by SSRO and Le Fort I osteotomy without orthodontic tooth extraction. At month 5 during 18 months of pre-surgical orthodontic treatment. The patient was completed after 7 months of post-surgical orthodontic treatment. The patient wore upper and lower Begg-type removable retainers for approximately 2 years. She returned for a recall checkup at 6 years post-treatment. Although radiographic examination revealed root resorption and

ankylosis of the autotransplanted tooth at 8 years after transplantation, occlusion has remained stable with no clinically significant complications. The autotransplanted tooth helped stabilize her occlusion and acted as a kind of temporary tooth prior to the final decision on treatment to be given such a dental implant.

Key words: Orthognathic surgery, Autotransplantation, Middle-aged patient, Long-term follow-up, Skeletal class III

[PDF (313K)] [References]

Download Meta of Article[<u>Help</u>] <u>RIS</u> BibTeX

To cite this article:

Etsuko Motegi, Yumi Takane, Eri Tokunaga, Kenji Sueishi, Nobuo Takano, Takahiko Shibahara and Chikara Saito: "Six-year Follow-up in Skeletal Class III Patient Aged over 40 Receiving Orthognathic Surgery and Autotransplantation: A Case Report". The Bulletin of Tokyo Dental College, Vol. **50**: 141-147 (2009).

doi:10.2209/tdcpublication.50.141 JOI JST.JSTAGE/tdcpublication/50.141

Copyright (c) 2009 by Tokyo Dental College, Japan

